

Citation for published version:

Li, Y & Qi, B 2017, 'Progress towards understanding protein S-acylation : prospective in plants', *Frontiers in Plant Science*, vol. 8, 346. <https://doi.org/10.3389/fpls.2017.00346>

DOI:

[10.3389/fpls.2017.00346](https://doi.org/10.3389/fpls.2017.00346)

Publication date:

2017

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link to publication](#)

Publisher Rights

CC BY

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Progress towards Understanding Protein S-acylation: Prospective in Plants

Yaxiao Li¹, Baoxiu Qi^{1*}

¹Biology and Biochemistry, University of Bath, United Kingdom

Submitted to Journal:
Frontiers in Plant Science

Specialty Section:
Plant Traffic and Transport

ISSN:
1664-462X

Article type:
Review Article

Received on:
14 Jan 2017

Accepted on:
28 Feb 2017

Provisional PDF published on:
28 Feb 2017

Frontiers website link:
www.frontiersin.org

Citation:
Li Y and Qi B (2017) Progress towards Understanding Protein S-acylation: Prospective in Plants. *Front. Plant Sci.* 8:346. doi:10.3389/fpls.2017.00346

Copyright statement:
© 2017 Li and Qi. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

Progress towards Understanding Protein S-acylation: Prospective in Plants

Yaxiao Li and Baoxiu Qi*

Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

*Correspondence: bssbq@bath.ac.uk

S-acylation, also known as S-palmitoylation or palmitoylation, is a reversible post-translational lipid modification in which long chain fatty acid, usually the 16-carbon palmitate, covalently attaches to a cysteine residue(s) throughout the protein via a thioester bond. It is involved in an array of important biological processes during growth and development, reproduction and stress responses in plant. S-acylation is a ubiquitous mechanism in eukaryotes catalyzed by a family of enzymes called Protein S-Acyl Transferases (PATs). Since the discovery of the first PAT in yeast in 2002 research in S-acylation has accelerated in the mammalian system and followed by in plant. However, it is still a difficult field to study due to the large number of PATs and even larger number of putative S-acylated substrate proteins they modify in each genome. This is coupled with drawbacks in the techniques used to study S-acylation, leading to the slower progress in this field compared to protein phosphorylation, for example. In this review we will summarize the discoveries made so far based on knowledge learnt from the characterization of protein S-acyltransferases and the S-acylated proteins, the interaction mechanisms between PAT and its specific substrate protein(s) in yeast and mammals. Research in protein S-acylation and PATs in plants will also be covered although this area is currently less well studied in yeast and mammalian systems.

Keywords: lipid modification, S-acylation, PATs, substrate recognition and specificity, yeast, mammalian, plants

INTRODUCTION

Lipid modification is a common mechanism in organisms, in which a fatty acid attaches to specific amino acid residues, leading to increased hydrophobicity of proteins which aids their anchoring to membranes or specific lipid rafts (Levental et al., 2010). The three most commonly known lipid modifications are N-myristoylation, prenylation and S-acylation (Figure 1). N-myristoylation is an irreversible, co-translational protein modification in which 14-carbon myristoyl group is covalently attached to N-terminal glycine residue via an amide bond (Martin et al., 2011). Prenylation is a post-translational lipid modification which involves the transfer of either a 15-carbon farnesyl or a 20-carbon geranyl-geranyl moiety to CaaX C-terminal cysteine of the target protein. S-acylation, more commonly known as S-palmitoylation, is a post-translational lipid modification in which a long chain fatty acid, usually the 16-carbon palmitate, covalently attaches to the specific cysteine residue(s) throughout the protein via a thioester bond (Resh, 2006; Greaves and Chamberlain, 2011).

It is noteworthy that three types of protein palmitoylation are found so far, including S-palmitoylation, N-palmitoylation and O-palmitoylation. While S-palmitoylation can occur at any Cys residues along the protein sequence in which the palmitate is reversibly attached via thioester bond as shown in Figure 1, N-palmitoylation is a stable lipid modification at the N-terminal residue (very often Cys) through amide linkage. A small group of secreted proteins have been identified as N-palmitoylated proteins, including the epidermal growth factor (EGF) like ligand 'Spitz' and Hedgehog family members in *Drosophila* and mammals (Pepinsky et al., 1998; Chamoun et al., 2001; Miura et al., 2006; Buglino and Resh, 2012). Since N-palmitoylation can be easily converted by S-palmitoyl migration, it is still not very clear whether N-palmitoylation is an independent enzyme-catalyzed reaction or just from S-to N-palmitoyl transfer (Ji et al., 2016). Less frequently, palmitoyl group can also be linked to a serine residue through ester bond via the so-called O-palmitoylation. The identified O-palmitoylated targets so far include Wnt/Wg proteins and the peptide

hormone preghrelin (Takada et al., 2006; Yang et al., 2008). Although palmitate is thought to be the most common fatty acid found to be attached to S-palmitoylated proteins recent studies proved that other acyl groups such as stearate (C18:0) or oleate (C18:1) are also accepted in S-palmitoylation. Therefore, S-acylation is a more representative term than palmitoylation (Jones et al., 1997; Sorek et al., 2007; Hurst and Hemsley, 2015). In contrast to other lipid modification, such as myristoylation, prenylation, N-palmitoylation or O-palmitoylation, S-acylation is a unique posttranslational modification in that it is usually reversible (Fukata et al., 2010). As such it is important for cellular protein sorting, vesicle trafficking, activation state control, protein stability, membrane microdomain partitioning of protein and protein complex assembly (Baekkeskov and Kanaani, 2009; Greaves and Chamberlain, 2007; Charollais and Van Der Goot, 2009; Hemsley, 2009; Hemsley et al., 2013).

Some other lipid modifications, such as Glycosylphosphatidylinositol (GPI) and glycosylinositolphosphorylceramide (GIPC) anchors can link the whole glycolipids to the protein instead of the simple fatty acid or polyisoprene group (Hemsley, 2015). GPI and GIPC anchors modify proteins at the lumen side instead of in the cytosol as do the other three lipid modifications (Ganesan and Levental, 2015). Lipid modifications which are only found in specific proteins, such as cholesterol addition at the C-terminal glycine of proteins have also been reported (Buglino and Resh, 2012). All these lipid modifications are widely present in mammals and plants (except for N- or O-palmitoylation which is only found in mammals so far) and all play important roles during growth and development through the modification of an array of proteins.

Although all lipid modifications can facilitate the attachment of proteins to membranes, modification with palmitoyl groups provide more affinity, about 10 times stronger than myristoyl groups and 100 times than farnesyl groups (Silvius and Heureux, 1994; Hemsley, 2009).

S-ACYLATION

S-acylation can occur both on soluble and transmembrane proteins (Roth et al., 2006; Blaskovic et al., 2013). S-acylation of soluble proteins allows their association with membranes, trafficking, regulation and signalling (Roth et al., 2006; Blaskovic et al., 2013). For example, a constitutive de/re-acylation of H- and N- small Rat sarcoma (Ras) drives their subcellular localization from plasma membrane (PM) to Golgi which initiates RAS activation (Rocks et al., 2005). Although the direct mechanism of S-acylation on transmembrane proteins is not very clear it is thought that it plays multiple roles in altering signalling capacity (Merrick et al., 2011), reducing activity (Huang et al., 2010), trafficking modification (Abrami et al., 2008; Flannery et al., 2010) and changing stability of these proteins (Maeda et al., 2010; Abrami et al., 2006; Blaskovic et al., 2013). For example, S-acylation of transmembrane proteins, such as death receptor 4 (Oh et al., 2012), β -secretase BACE1 (Motoki et al., 2012), cannabinoid receptor (Oddi et al., 2012) and influenza virus M2 protein (Thaa et al., 2011), can promote their association with membrane lipid rafts. However, for some peripheral membrane proteins such as transferrin receptor and caveolin, their palmitoylation sites are localized to non-raft domains, therefore palmitoylation is not necessary for their raft localization (Alvarez et al., 1990; Dietzen et al., 1995; Charollais and Van Der Goot, 2009). In the case of the tumor endothelial marker 8 (TEM8) palmitoylation was actually found to negatively regulate its raft association (Abrami et al., 2006).

S-acylation in Yeast

A proteomic method using the acyl-biotinyl exchange (ABE) chemistry combining with the traditional [^3H] palmitate in vivo labeling protocol identified 48 S-acylated proteins that span a wide range of cellular functions in *Saccharomyces cerevisiae* (Roth et al., 2006). These include a large number of SNAREs (soluble N-ethylmaleimide-sensitive fusion protein-attachment protein receptor) that are involved in vesicle fusion. Redundant SNAREs, such as plasma membrane (PM) localized synaptobrevin homologs Snc1 and Snc2, were first identified to be S-acylated proteins in 1995 (Couve

et al., 1995), and subsequently confirmed independently (Valdez-Taubas and Pelham, 2005; Roth et al., 2006). Ykt6 is another commonly known S-acylated SNARE. It requires both C-terminal prenylation and palmitoylation to target to the membrane, which is different from all other single transmembrane domain (TMD) containing SNAREs (Fukasawa et al., 2004). Tlg1 lacking S-acylation undergoes ubiquitination, implying S-acylation can protect proteins from degradation (Valdez-Taubas and Pelham, 2005). Other SNAREs that have been confirmed to be S-acylated are Sso1, Sso2, Vam3, Tlg2 and Syn8 (Valdez-Taubas and Pelham, 2005; Roth et al., 2006). S-acylation is also very common in many important signalling proteins, such as the heterotrimeric G protein alpha and gamma subunits Gpa1 (Song and Dohlman, 1996; Song et al., 1996), Gpa2 (Harashima and Heitman, 2005) and Gγ (Ste18, Hirschman and Jenness, 1999); small monomeric G proteins (GTPases) such as Rho1, Rho2 (Roth et al., 2006), Rho3 (Zhang et al., 2013), Ras1 and Ras2 (Deschenes et al., 1990; Bartels et al., 1999; Mitchell et al., 2012). A recent study shows that the pathogenesis, morphogenesis and sexual differentiation of an encapsulated yeast *Cryptococcus neoformans* is achieved through the important roles that S-acylation plays in modulating the localization of Ras1 (Nichols et al., 2015). Interestingly, all of these signaling proteins acquire prenylation or myristoylation before S-acylation occurs (Roth et al., 2006).

In addition, many amino acid permeases (AAP) were proved to be S-acylated (Roth et al., 2006). For example, the yeast type I casein kinases, Yck1, Yck2, and Yck3, which play important roles in cellular morphology, bud emergence and endocytosis of mating pheromone receptor, are membrane localized via S-acylation for function (Roth et al., 2006; Roth et al., 2011). ENV7 (late endosome and vacuole interface) encodes a protein kinase that plays important roles in vacuole morphology, and its proper membrane localization and function relies on S-acylation of the N-terminal triple cysteines motif (C¹³C¹⁴C¹⁵) (Manandhar et al., 2013 and 2014; Cocca, 2014). S-acylation of telomere-binding protein Rif1 anchored it to the inner nuclear membrane, which influences its role in heterochromatin dynamics (Park et al., 2011). Mutagenesis of cysteine in different positions of Arsenite permease Acr3p can cause its completely or partially

dysfunction as a low affinity As(III)/H⁺ and Sb(III)/H⁺ antiporter, and Cys90 which localizes in the cytosolic loop but in close proximity to transmembrane regions has the high possibility to be S-acylated (Maciaszczyk-Dziubinska et al., 2013). It was also reported that S-acylation is necessary for the export of chitin synthase Chs3 from ER (Lam et al., 2006). The information described in this section is summarized in **Table 1**.

TABLE 1| Individually confirmed S-acylated proteins in yeast

Groups	Specific proteins	References
SNAREs	Snc1/2, Ykt6, Tlg1/2, Sso1/2, Vam3, Syn8	Couve et al., 1995; Valdez-Taubas and Pelham, 2005; Roth et al., 2006; Fukasawa et al., 2004
G proteins	Gpa1/2, Ste18, Rho1/2/3, Ras1/2	Song and Dohlman, 1996; Song et al., 1996; Hirschman and Jenness, 1999; Harashima and Heitman, 2005; Roth et al., 2006; Zhang et al., 2013; Deschenes et al., 1990; Bartels et al., 1999; Mitchell et al., 2012; Nichols et al., 2015
AAPs	Tat1/2, Gnp1, Sam3, Hip1, Bap2, Agp1, Gap1	Roth et al., 2006
Protein kinases	Yck1/2/3, Env7	Roth et al., 2006; Roth et al., 2011; Cocca, 2014
Other proteins	Rif1, Acr3p, Chs3	Park et al., 2011; Maciaszczyk-Dziubinska et al., 2013; Lam et al., 2006

S-acylation in Mammals

Following study of S-acylation in yeast research that has extended to mammalian systems considerable knowledge has been gained in recent years, revealing the involvement of protein S-acylation in the regulation of growth, development, and cancer and disease status. For example, a global rat neural palmitoyl-proteome

characterized almost 300 S-acylated proteins, again with the ABE method adapted from the yeast study (Kang et al., 2008). Similarly 331 S-acylated proteins were identified from human prostate cancer cells (Yang et al., 2010), 57 from human B lymphoid cells (Ivaldi et al., 2012) and 150 from endothelial cells (Marin et al., 2012). By bio-orthogonal labeling of S-acylated proteins with 17-octadecynoic acid (ODYA) about 125 and over 400 S-acylated proteins were identified from human Jurkat T-cells and mouse T-cell hybridoma cells, respectively (Martin and Cravatt, 2009; Martin et al., 2012).

It is worth noting that proteins that have been proved to be S-acylated in yeast, their homologous proteins in mammals tend to be also S-acylated. For instance, many human SNAREs were proved to be also S-acylated (Greaves et al., 2010), S-acylation of α subunits of G proteins is necessary for their membrane localization and function (Wedegaertner et al., 1993; Grassie et al., 1994; Ponimaskin et al., 1998; Ponimaskin et al., 2000). However, G-protein γ subunits have not been reported to be S-acylated in mammals. Many G-protein-coupled receptors (GPCRs) (Blaskovic et al., 2013) and Ras GTPase (Rocks et al., 2005) are also S-acylated. Mitochondrial targeting of a microphage protein phospholipid scramblase 3 (Plscr3) is dependent on its S-acylation (Merrick et al., 2011).

Some S-acylated proteins in mammals can also make themselves avoid degradation by attaching a palmitate molecule. For instance, LRP6 (lipoprotein-receptor-related protein 6) is S-acylated and the removal of acyl group leads to destabilization or ubiquitination (Abrami et al., 2008). Similarly, the palmitoylation of TEM8 (Abrami et al., 2006), CCR5 (chemokine and HIV receptor) (Percherancier et al., 2001) and Rhodopsins (Maeda et al., 2010) prevents the degradation of these proteins. It was also reported that for some other proteins, their degradation depends on the S-acylation. For example, a cancer-promoting protein CDCP1 (CUB domain-containing protein 1) is degraded upon S-acylation, leading to a decrease of ovarian cancer cell migration (Adams et al., 2015). Therefore, it seems that S-acylation can play opposite roles in

protein degradation.

Many signalling proteins involved in keeping T-cell homeostasis are S-acylated, such as T-cell co-receptors CD4 and CD8, tyrosine kinases Lck and Fyn, and adaptor proteins LAT (linker for activation of T cells) and Cbp/PAG (Bijlmakers, 2009; Hundt et al., 2009; Akimzhanov and Boehning, 2015). S-acylation of Lck at both Cys3 and Cys5, which are redundant for the function of Lck, is essential for propagating T-cell receptor signaling and releasing apoptotic calcium (Akimzhanov and Boehning, 2015). Similarly, LAT is also a dual (Cys26 and Cys29) S-acylated protein which is required for T cell development and activation. However, S-acylation of Cys26 alone is enough for its PM localization and proper function (Hundt et al., 2009).

S-acylation of synaptic proteins is important for synaptic plasticity, and the key S-acylated synaptic proteins include postsynaptic density protein PSD-95, δ -catenin, gephyrin, A-kinase anchoring protein AKAP79 and 150, the small GTPase Cdc42. Lack of S-acylation of these proteins lead to impaired performance on learning and memory tasks (Brigidi et al., 2015). Huntington's disease is a neurodegenerative disorder caused by mutation in the gene encoding the S-acylated Huntingtin (HTT) (Butland et al., 2014). Defects in S-acylation can also cause mental problems such as schizophrenia and X-linked mental retardation (XLMR), however, the specific S-acylated target proteins involved in this process have not been isolated (Mukai et al., 2004; Raymond et al., 2007). Alzheimer's disease (AD) is a neurodegenerative dementia which accounts for 60% to 70% of cases of dementia. Many studies have demonstrated that S-acylation plays very important roles in the pathogenesis of AD, and the related S-acylated proteins include β - and γ -secretase enzymes, and the major APP (amyloid precursor protein) cleaving enzyme BACE1, which are S-acylated at four sites (Benjannet et al 2001; Hornemann, 2015).

Autophagic protein microtubule-associated protein 1 light chain-3B (LC3B) is a positive regulator of chronic obstructive pulmonary diseases such as emphysema. LC3B is associated with the extrinsic apoptotic factor Fas, and their interaction is

mediated by caveolin-1 (Cav-1). Interestingly, both Fas and Cav-1 are S-acylated proteins (Chen et al., 2010). S-acylation of the bone developmental regulator membrane type1-metalloprotease (MT1-MMP) is a key modulator of bone homeostasis (Song et al., 2014). Goltz syndrome, caused by loss of function of the S-acylated protein Porcupine (Galli et al., 2007; Hornemann, 2015), is an X-linked dominant form of ectodermal dysplasia, which is primarily characterized by skin manifestations as atrophic and hypoplastic areas and results in osseous defects and dental anomalies later (Wang et al., 2007).

An increasing number of reports indicate that S-acylation is involved in cancer. For instance, Ras is a negative regulator of cell proliferation, and S-acylation of Ras maintains its steady state plasma membrane localization which is essential for transduction of extracellular proliferative signals (Rocks et al., 2005; Schmick et al., 2015). S-acylation of the neurotensin receptor 1 (NTSR-1), a key mediator in breast, pancreas, prostate, colon and lung cancers, is essential for its localization and efficient signaling (Heakal et al., 2011). The induction of apoptosis is an efficient way to stop tumor development, many proteins involved in apoptosis are S-acylated including FasL (Fas Ligand) (Guardiola-Serrano et al., 2010), FasR (Fas receptor) (Chakrabandhu et al., 2007), DR4 (a receptor of the tumor necrosis factor-related apoptosis-inducing ligand) (Rossin et al., 2009), DCC (deleted in colorectal cancer) (Furne et al., 2006), UNC5H (Maisse et al., 2008) and BAX (BCL-2-associated X) (Fröhlich et al., 2014). The spread of cancer cells from their original site to other parts of the body is through metastasis. It was reported that S-acylation of metastasis-associated proteins KAT1/CD82, CD9 and CD151 is essential for their function of suppressing metastasis or inhibiting tumor cell adhesion and migration (Zhou et al., 2004; Termini et al., 2014; Hemler, 2014). Integrin $\beta 4$ (ITG $\beta 4$) can interact with growth factor receptors and enhance invasive potential of cancer cells (Soung and Chung, 2011). This is helped by the S-acylation of ITG $\beta 4$ which is required for its lipid raft localization in the membrane and signaling activity. The level of ITG $\beta 4$ S-acylation is correlated with the invasive potential of breast cancer cells (Coleman et al., 2015). Another S-acylated protein

related to breast cancer is CD44 which negatively regulates cell migration (Xie et al., 2009). Endothelial nitric oxide synthase (eNOS), which localizes through S-acylation to the Golgi complex and PM cholesterol-rich microdomains, promotes angiogenesis and tumorigenesis (Fernandez-Hernand et al., 2006; Wei et al., 2011). **Table 2** lists the identified S-acylated proteins in mammals described in this section.

TABLE 2 | S-acylated proteins individually verified in mammalian cells

Groups	Examples	References
SNAREs	SNAP23, SNAP25, SNAP25b	Greaves and Chamberlain, 2010; Greaves et al., 2010
G Proteins	Go1 α , G α ₁₂ , G α ₁₃ , GPCRs, GTPase	Wedegaertner et al., 1993; Grassie et al., 1994; Ponimaskin et al., 1998; Ponimaskin et al., 2000; Rocks et al., 2005
T-cell specific proteins	CD4/8, Lck, Fyn, LAT, Cbp/PAG	Bijlmakers, 2009; Hundt et al., 2009; Akimzhanov and Boehning, 2015
B-cell specific proteins	CD20/23	Ivaldi et al., 2012
Synaptic proteins	PSD-95, δ -catenin, gephyrin, AKAP79/150, Cdc42, HTT, β - and γ - secretases, BACE1	Brigidi et al., 2015; Butland et al., 2014; Benjannet et al 2001; Hornemann, 2015
Cancer related proteins	CDCP1, Ras, NTSR-1, FasL, FasR, DR4, DCC, UNC5H, BAX, CD82/9/151/44, ITG β 4, Enos	Adams et al., 2015; Schmick et al., 2015; Heakal et al., 2011; Guardiola-Serrano et al., 2010; Chakrabandhu et al., 2007; Rossin et al., 2009; Furne et al., 2006; Maisse et al., 2008; Fröhlich et al., 2014; Zhou et al., 2004; Termini et al., 2014; Hemler, 2014; Soung and Chung, 2011; Coleman et al., 2015; Xie et al., 2009; Fernandez-Hernand et al., 2006; Wei et al., 2011

Other Proteins	Plscr3, LRP6, Fas, Cav-1, MT1-MMP; Porcupine, TEM8, CCR5	Merrick et al., 2011; Abrami et al., 2008; Abrami et al., 2006; Percherancier et al., 2001; Chen et al., 2010; Song et al., 2014; Galli et al., 2007
----------------	--	--

245

246 The above studies clearly demonstrate that S-acylation is involved in a wide range
247 of human diseases including mal-development, infectious diseases, autoimmune
248 diseases, neuropsychiatric disorders, dermatosis, osteoporosis and cancer (Ivaldi et al.,
249 2012; Chavda et al., 2014; Hornemann, 2015; Yeste-velasco et al., 2015).
250 Understanding S-acylation will provide invaluable information to the insight of disease
251 processes which in turn will aid the development of drugs to control and target these
252 various diseases. Therefore, the relationship between protein S-acylation and disease in
253 human becomes a hot research topic in the medical field in recent years.

254 **S-acylation in Plants**

255 Our understanding of plant S-acylation is rudimentary and the limited knowledge
256 comes mainly from targeted studies on the functional characterization of individual
257 proteins that happen to be S-acylated, including, mainly heterotrimeric G protein and
258 some small monomeric G-proteins. For instance, the α subunit GPA1 and γ subunit
259 AGG2 of plant heterotrimeric G protein are S-acylated. GPA1 has dual lipid
260 modification with a myristoylation site at the G2 position and an adjacent S-acylation
261 site at the C5 position, ensuring its localization to the PM (Adjobo-Hermans et al.,
262 2006). Apart from promoting PM localization, S-acylation of GPA1 may also stabilise
263 the newly formed heterotrimer. AGG2 is S-acylated at Golgi before delivered to the PM,
264 and its membrane localization is dependent on its prenylation and S-acylation (Zeng et
265 al., 2007). Therefore, S-acylation may act as a membrane targeting signal and restricts
266 AGG2 shuttle in and out of PM (Zeng et al., 2007; Hemsley, 2009). Some small
267 GTPases are also known to be S-acylated. For instance, S-acylation of AtROP6 is

responsible for its activation and inactivation cycles (Sorek et al., 2007). AtROP9 and AtROP10, which are involved in ABA signalling, contain 3 and 2 S-acylation sites respectively (Lavy et al., 2002; Zheng et al., 2002; Hemsley, 2009). For AtRABF1 (ARA6), both S-acylation and myristoylation are essential for its prevacuolar compartment localization (Ueda et al., 2001).

Proteins involved in Ca^{2+} signalling such as calcineurin B-Like proteins AtCBL1, AtCBL2, AtCBL3 and AtCBL6 in Arabidopsis (Batistic et al., 2008; Batistic et al., 2012); calcium dependent protein kinases OsCPK2 in rice (Martin and Busconi, 2000); LeCPK1 in tomato (Leclercq et al., 2005); MtCPK3 in *Medicago truncatula* (Gargantini et al., 2006) and StCDPK1 in *Solanum tuberosum* (Raices et al., 2003) were reported to be S-acylated. AtCBL1 is a dually lipid modified protein, in which myristoylation targets it to the endoplasmic reticulum (ER), but the trafficking from ER to PM and subsequent PM anchoring depends on S-acylation (Batistic et al., 2008).

Other S-acylated proteins are the pathogenesis related proteins such as RPM1 interacting protein 4 (RIN4) and leucine-rich repeat receptor like kinase (FLS2) (Kim et al., 2005; Hemsley et al., 2013; Running, 2014; Boyle et al., 2016); NDR1/HIN1-like (NHL) stress response proteins (Hemsley et al., 2013; Hurst and Hemsley, 2015); POLTERGEIST (POL) and POLTERGEIST LIKE 1 (PLL1) (their PM localization is dependent on both myristoylation and S-acylation at their N-termini) (Gagne and Clark, 2010); the Lost In Pollen tube guidance 1 (LIP1) and 2 (LIP2), mutations of their S-acylation sites abolished PM localization. Although individual knockout mutant of LIP1 and LIP2 did not have any defects the double mutant can cause sterility due to loss of pollen tube guidance (Liu et al., 2013). S-acylation of remorin proteins, a group of well-known plasma membrane marker proteins, contribute to their subcellular localization (Konrad et al., 2014). Very recently, Kumar and his coworkers confirmed that a number of the catalytic subunits of cellulose synthase complex (CSC) in Arabidopsis are S-acylated. These include the cellulose synthase A 1 (CESA1), CESA4, CESA6, CESA7 and CESA8 where up to 6 S-acylation sites were in each of these

proteins (Kumar et al., 2016). SGN1, a receptor-like cytoplasmic kinase (RLCK), localizes in a strictly polar fashion to the endodermal outer plasma membrane, and this is dependent on the S-acylation of N-termini (Alassimone et al., 2016) (See summary in Table 3).

TABLE 3 | S-acylated proteins individually identified in plants

Groups	Examples	References
SNAREs	AtSYP71, AtSYP122, AtNPSN11	Hemsley et al., 2013
G-proteins	AtGPA1, AtAGG2, AtROP6/9/10, AtRABF1	Ueda et al., 2001; Lavy et al., 2002; Zheng et al., 2002; Adjobo-Hermans et al., 2006; Sorek et al., 2007; Zeng et al., 2007
Proteins in Ca ²⁺ signalling	AtCBL1/2/3/6, OsCPK2, LeCPK1, MtCPK3, StCDPK1	Martin and Busconi, 2000; Raices et al., 2003; Leclercq et al., 2005; Gargantini et al., 2006; Batistic et al., 2008; Batistic et al., 2012;
Cellulose Synthase complex	AtCESA1, AtCESA4, AtCESA6, AtCESA7, AtCESA8	Kumar et al., 2016
Others	RIN4, FLS2, POL, PLL1, LIP1, LIP2, remorins, SGN1	Kim et al., 2005; Gagne and Clark, 2010; Hemsley et al., 2013; Running, 2014; Liu et al., 2013; Konrad et al., 2014; Alassimone et al., 2016

On a proteomic level Hemsley and coworkers identified about 600 putative S-acylated proteins from Arabidopsis using a biotin switch isobaric tagging for relative and absolute quantification (Hemsley et al., 2013). These proteins are involved in many processes across plant growth, development and stress responses, including the

mitogen-activated protein kinases (MAPKs), leucine-rich repeat receptor-like kinases (LRR-RLKs) and RLK superfamily members, integral membrane transporters, ATPases, SNAREs and others. Similarly, about 450 S-acylated proteins were identified from Poplar cell suspension very recently. Except for the commonly known intracellular trafficking related proteins such as protein kinases, SNAREs, band 7 family proteins and tetraspanins, some cell wall related proteins were also found to be S-acylated (Srivastava et al., 2016). These results greatly expand the range of functions of protein S-acylation involves in plants, demonstrating the important roles of protein S-acylation in plant growth, development and stress signaling.

S-acylation in Other Organisms

S-acylated proteins were also identified from other organisms. For example, more than 400 putative S-acylated proteins were isolated from the most severe human malaria causing parasite *Plasmodium falciparum*, involved in almost all the stages of its life cycle (Hodson et al., 2015). A number of S-acylated proteins are localized in the inner membrane complex (IMC). IMC is a membranous two layered structure located underneath the plasma membrane, which IMC plays central roles in host cell invasion and cytokinesis in *Plasmodium falciparum* (Cavalier-Smith, 1993; Wetzel et al., 2014). In another parasite *Toxoplasma gondii*, S-acylated proteins were also proved to be involved in many physiological processes including motility, invasion and division (Frenal et al., 2010; Beck et al., 2010; Frenal et al., 2014). In *Aspergillus fumigatus*, one of the most common species that cause the invasive aspergillosis in individuals with an immunodeficiency, Ras pathway signaling is its critical virulence determinant and the properly localized and activated Ras is dependent on a series of posttranslational lipid modification including S-acylation (Abdallah and Fortwendel, 2005). Study also showed that S-acylation is essential for spermatogenesis of *Caenorhabditis elegans* (Gleason et al., 2006).

S-acylation not only occurs on proteins synthesized in eukaryotic cells but also for proteins secreted by prokaryotic bacteria and viruses and subsequently S-acylated by

their eukaryotic hosts. For example, *Legionella* and other bacterial pathogens can secrete effectors that mimic the substrates of host lipid transferases, which can help them target the proper host membranes after S-acylation and other lipid modifications (Ivanov and Roy, 2013). A group of cysteine protease type III effectors secreted by the plant pathogen *Pseudomonas syringae*, rely on their S-acylation by the host cells to be targeted to plasma membrane and activated (Downen et al., 2009). S-acylation can also take place on viral proteins and in fact the first reported S-acylated protein was a glycoprotein from *Vesicular stomatitis* virus (Schmidt and Schlesinger, 1979; Hurst and Hemsley, 2015). Another viral S-acylated protein is the hemagglutinin of Influenza virus. S-acylation of its all three cysteine residues by the host cell S-acylation machinery is essential for the replication and infection of the virus (Zurcher et al., 1994; Wagner et al., 2005; Brett et al., 2014).

PROTEIN S-ACYL TRANSFERASES (PATs)

While spontaneous palmitoylation does occur on some proteins in the cells (i.e. Bizzozero et al., 2000; Kummel et al., 2006; Kostiuk et al., 2008) it is generally accepted that S-acylation is an enzymatic process catalyzed by a family of proteins, the Protein S-Acyl Transferases (PATs for short). This is because research on PATs was much delayed compared to that on the S-acylated proteins. The first PAT, Akr1 was identified from *Saccharomyces cerevisiae* in 2002 which is 20 years later than S-acylation of protein first reported (Roth et al., 2002; Schmidt and Schlesinger, 1979; Bartels et al., 1999). Since then, the significance of this enzyme family has been gradually recognized by studies carried out by an increasing number of researchers in this field, leading to the great enrichment of our knowledge of PATs, particularly in yeast and mammals.

The Structure and Functional Domains of PATs

Compared to the numbers of enzymes that catalyze the N-myristoylation or prenylation, there are much more DHHC-containing PATs existing in eukaryotes. In contrast to the

cytoplasmic catalyzing enzymes for S-prenylation and N-myristoylation, PATs are transmembrane proteins with 4-6 TMDs and cytosolic N- and C- termini (Hemsley et al., 2013). Most importantly, PATs also have a highly conserved catalytic Asp-His-His-Cys Cysteine Rich Domain (DHHC-CRD) of ~50 amino acids (Roth, et al., 2002). This domain was proposed as Cx2Cx9HCx2Cx4DHHCx5Cx4Nx3F (Mitchell et al., 2006), usually residing on the cytoplasmic face of membranes between transmembrane domains (TMD) 2 and 3 of PATs (Gottlieb et al., 2015). It was reported that mutation of cysteine in DHHC domain inhibits both acyl intermediate formation and acyl chain transfer activity of PATs (Mitchell et al., 2006; Gottlieb et al., 2015). Indeed, when cysteine residue in the DHHC motif of AtPAT24, AtPAT10 and AtPAT14 of Arabidopsis was mutated to alanine or serine, all 3 AtPATs lost their PAT activities (Hemsley et al., 2005; Qi et al., 2013; Li et al., 2015). The DHHC-CRD domain in Swf1 cannot be replaced by those from Pfa3, Pfa4 or Erf2, and similar results were also found for Pfa3, the DHHC-CRD of which cannot be replaced by that of Swf1 or Erf2. The irreplaceability of DHHC-CRD demonstrates interaction between this domain and other regions is required for proper PAT function (Gonzalez Montoro et al., 2011). Although the acyl intermediate happened on the cysteine residue in the DHHC motif study on human DHHC3 showed that mutation of other conserved cysteines in the CRD also decreased its activity (Gottlieb et al., 2015). In addition, cysteine residues within a novel CCX₇₋₁₃C(S/T) motif downstream of the conserved DHHC-CRD of human PATs DHHC5, DHHC6 and DHHC8 were also proved to be S-acylated (Yang et al., 2010). Therefore, it seems that cysteine residues in the DHHC-CRD as well as other motifs play joint roles in of PATs auto-acylation and subsequent transfer of the fatty acid to their substrate proteins. It is also interesting to note that many residues in the DHHC-CRD domain are reported to determine substrate specificity of a PAT, such as A145 and K148 in Swf1 (Gonzalez Montoro et al., 2011).

Some PATs also have an N-terminal ankyrin repeat (AR) domain. Usually two AR containing DHHC PATs are found in per genome, such as in mammals (Fukata et al., 2004), yeast (Roth et al., 2006), fly (Bannan et al., 2008), apicomplexan parasite (Frenal

et al., 2013), nematode (Edmonds and Morgan, 2014) and plants (Yuan et al, 2013). It is thought that AR can help these PAT to recognize its specific targets for S-acylation (Lemonidis et al., 2015). However, other functions of AR that are independent from S-acylation were also found in some such PATs (Harada et al., 2003; Yang and Cynader, 2011; Hemsley and Grierson, 2011). For example, AR is essential for Akr1 to interact with G β γ dimer and to suppress cell cycle arrest induced by β subunit, and this process does not require Akr1 being a functional PAT (Hemsley and Grierson, 2011).

Both the N- and C-termini of all PATs characterized so far are highly variable and cytosolic. The highly variability of N- and C- terminal domains are believed to be essential for substrate specificity of PATs, even though there have no experimental evidence to support this at present (Huang et al., 2009; Greaves et al., 2010; Gonzalez Montoro et al., 2011). Many PATs also have a conserved aspartate-proline-glycine (DPG) motif close to the second TMD, and a threonine-threonine-asparagine-glutamate (TTxE) motif adjacent to the last TMD. Both of them are cytosolic but their roles in the function of PATs are still waiting to be explored. Another important region that contains 16-amino acids and is conserved in 70% eukaryotic PATs is the PaCCT motif (Palmitoyltransferase Conserved C-Terminus). Absence of the PaCCT motif abolished the function of Pfa3 in yeast; the tyrosine residue within this motif of Swf1 is essential for its PAT activity towards Tlg1 (Gonzalez- Montoro et al., 2009).

The general structure and the functional/conserved domains of PATs is illustrated in **Figure 2**.

DHHC Proteins Are Commonly Found in Eukaryotes

Since the first DHHC containing protein, Akr1 was found and proved to be a PAT from yeast in 2002, significant advances have been made in understanding of DHHC protein family in yeast, mammals, worm and plants. So far, 6 of the 7 yeast DHHC proteins have been confirmed to be PATs, and they are Akr1 (Roth et al., 2002; Lobo et al., 2002), Erf2 (Valdez-Taudas and Pelham, 2005), Swf1 (Smotrys et al., 2005), Pfa3 (Hou et al.,

2009), Pfa4 (Smotrys et al., 2005) and Pfa5 (Roth et al., 2006). Akr2 is highly homologous to Akr1 with a typical DHHC-CRD and two ARs, however, *akr2* mutant did not show any remarkable phenotype and there is no direct evidence to show whether it has PAT catalytic activity or not (Kihara et al., 2005; Linder and Deschenes, 2007).

Among the 22 human DHHC proteins (DHHC1-22), 17 were proved to have PAT activities excluding DHHC4, 11, 13, 19 and 22 (Ohno et al., 2012). One more DHHC protein (DHHC23) was found in mice but its homologue cannot be found in human (Ohno et al., 2006; Greaves and Chamberlain, 2010). *Caenorhabditis elegans* has 15 DHHC-PATs, but so far only 1, SPE10 (spermatogenesis 10), was characterized in some detail and showed that it is essential for membranous organelles to deliver fibrous bodies to the spermatid (Gleason et al., 2006).

Plant genomes also possess various numbers of DHHC containing protein sequences. A recent survey from 31 plant species with complete genomes including Arabidopsis, identified 804 DHHC proteins. The numbers of DHHC proteins were variable in different species from 6 in *volvox carteri* to 52 in *Panicum virgatum*. Expression pattern of DHHC proteins in *Zea mays* and their response to phytohormones and abiotic stress showed that these DHHC proteins may play important roles in plant growth and development as well as stress responses (Yuan et al., 2013). Arabidopsis has 24 DHHC-containing proteins, named as ATPAT1-24 (Hemsley et al, 2005; Batistic, 2012). According to their phylogenetic relationship they are divided into 3 main groups, where group A has the most members including AtPATs 1 to 9, group B consists AtPATs 11-16, group C is made of AtPATs 18 to 22, whilst AtPAT10, 17, 23 and 24 do not belong to any groups (Batistic, 2012). All these putative PATs have 4 TMDs except for AtPAT15 and AtPAT17 where 3 and 6 TMDs are found respectively.

Similar to what was found in yeast and mammals, Arabidopsis genome also has 2 ankyrin repeats containing PATs, AtPAT23 and AtPAT24 that are highly homologous. Being the first PAT identified in higher plant, AtPAT24 was confirmed to be an S-acyl transferase because it not only auto-acylated but also rescued the yeast PAT Akr1

knockout mutant *akr1* for its morphological and temperature sensitive defects. AtPAT24 can also restore the correct localization of one of the Akr1 palmitoylating proteins, the yeast casein kinase 2 (Yck2). The AtPAT24 loss-of-function mutant *tip1* exhibits defects in cell size control, pollen tube and root hair growth, as well as cell polarity (Hemsley et al., 2005). Following this, the biological functions of 3 other AtPATs have also been characterized in some detail recently. For example, three T-DNA insertion mutant alleles were identified for AtPAT10 and all showed pleiotropic defects, including cell expansion, cell division, vascular patterning, fertility and salt stress in Arabidopsis (Qi et al., 2013; Zhou et al., 2013). Single mutants of *atpat13* and *atpat14* are semi-dwarf and show precocious leaf senescence, and the double mutant *atpat13 atpat14* has even stronger phenotype than each of their parent single mutant plants (Lai et al., 2015; Li et al., 2015).

DHHC containing proteins were also identified from other organisms. For example, 22 such proteins were found in *Drosophila* (Bannan et al., 2008), 18 in *Toxoplasma gondii*, 17 in *Neospora caninum*, 12 in *Trypanosoma brucei*, 12 in *Plasmodium falciparum*, 11 in *Plasmodium berghei*, 10 in *Cryptosporidium* species, 9 in *Theileria parva*, 8 in *Babesia bovis* and 6 in *Eimeria tenella* (Frénal et al., 2013). It was reported that TgDHHC7 from *Toxoplasma gondii* is essential for rhoptry organelles localization and parasite invasion (Frénal et al., 2013).

Expression Pattern and Subcellular Localization of PATs

The spatial and temporal expression patterns of genes are very important for their cellular functions. However, only very limited information available for the expression patterns of PAT family proteins so far. It was shown that the humans DHHC1, 3-10, 12-14, 16-18 and 20-22, are ubiquitously expressed in different tissues. DHHC19 had very high expression level in testis with weak expression in thymus and small intestine while DHHC11 was only expressed in testis. In addition, only very low level of *DHHC2* transcript was present in kidney and testis, and similar low levels of *DHHC15* transcript were found in heart, brain, lung, kidney, thymus, and small intestine (Ohno et al., 2006).

The expression patterns of DHHC proteins in *Drosophila* were also analyzed. Among them, CG1407, CG5620, CG6017, CG6627 and CG17257 exhibited maternal expression and were enriched in neural tissues, with transcripts of all of them except for CG1407 also detected in larval brains. Some DHHC proteins were only expressed in testis, such as CG4483, CG4956, CG13029, CG17075, CG17195-17198, CG17287 and CG18810, and others expressed only in ovary, such as CG5880, CG6017 and CG34449 (Bannan et al., 2008).

Study of the expression patterns of PATs in parasite also showed that most of them have ubiquitous expression with a few being more tissue or developmental stage specific. For instance, TgDHHC18 is specially expressed in bradyzoites, TgDHHC10 at oocyst stage, PfDHHC6 and PfDHHC10 at gametocyte stage (Frenal et al., 2013; Lopez-Barragan et al., 2011).

In *Arabidopsis*, 19 of the 24 *AtPATs* expressed in a broad and constant pattern with transcripts detected in most tissues at all different developmental stages in *Arabidopsis*. However, *AtPAT1*, 2, 3, 11 and 21 had relatively low expression levels than other *AtPATs*, and *AtPAT2* and 3 also exhibited stronger expression in pollen (Batistič, 2012; Yuan et al., 2013). In *Oryza sativa*, 26 of the 30 *OsPATs* can express in more than one type of tissue, among which *OsPAT29* was only expressed during germination stage, *OsPAT21* and *OsPAT26* were only expressed in the internode and stamen respectively. The transcripts of *OsPAT13* and *OsPAT28* were barely detectable in the tissues examined (Yuan et al., 2013). In *Zea mays* 28 of 38 *ZmPATs* have extensive expression in different developmental stages and tissues with *ZmPAT13* and *ZmPAT22* having higher expression in anther (Yuan et al., 2013). However, in *Glycine max*, specific, rather than broad expression patterns were found where 7 *GmPATs* were specifically expressed in the flowering stage and the transcripts of the remaining 11 *GmPATs* were detected in all developmental stages except for flowering stage (Yuan et al., 2013).

Therefore, it is clear that most PATs exhibit a broad expression pattern in different developmental stages and tissues with a few stage or tissue specific PATs in all

organisms reported. This suggests that most PATs are involved in a broad range of functions in a given organism, such as Arabidopsis.

PATs distributed in the entire endomembrane system in the cell and this locality nature of PATs may determine the specific set of proteins they modify. For example, the plasma membrane-localized Pfa5 in yeast and DHHC5, 20 and 21 in human are the PATs involved in S-acylation mediated signal transduction of PM localized heterotrimeric G protein alpha subunit G α , the β 2-adrenergic receptor and endothelial nitric oxide synthase (Mumby et al., 1994; Loisel et al., 1996; Robinson et al., 1995; Ohno et al., 2006). The ER- and Golgi-localized DHHC proteins may be responsible for palmitoylation of de novo synthesized proteins during the processes of membrane localization and delivery to other organelles (Ohno et al., 2006). The tonoplast-localized Pfa3 in yeast palmitoylates Vac8p for its vacuolar membrane targeting (Hou et al., 2005; Smotrys et al., 2005). Therefore, to determine the subcellular localization of individual PAT is very important in order to understand its function by identifying the substrate protein(s) it modifies and signaling pathways it is involved.

Although PATs are found in all endomembrane systems in the eukaryotic cell they have different preference in different species as to which endomembrane compartment they are localized. For instance, in yeast, 3 PATs, Swf1, Pfa4 and Erf2 are localized at ER; Akr1 and Akr2 are localized at Golgi; while Pfa3 and Pfa5 are localized at vacuole and plasma membrane respectively (Valdez-Taubas and Pelham, 2005; Ohno et al., 2006). In human, 8 PATs are localized at ER (DHHC1, 6, 10, 11, 13, 14, 16, 19), 7 at Golgi (DHHC3, 4, 7, 8, 15, 17, 18), 2 at PM (DHHC5 and 20), 4 at both ER and Golgi (DHHC2, 9, 12 and 22), and 1 at both Golgi and PM (DHHC21) (Ohno et al., 2006).

Similar to the mammalian PATs, DHHC-PATs in *Drosophila* are also mainly localized at ER (14: CG4483, CG4676, CG4956, CG5196, CG5620, CG6627, CG10344, CG13029, CG17075, CG17195, CG17196, CG17197, CG17198 and CG17287) and Golgi (6: CG5880, CG6017, CG6618, CG8314, CG17257 and CG18810). The only exception is CG1407 which is localized at PM (Bannan et al.,

2008). In Apicomplexan, such as *Toxoplasma gondii*, TgDHHCs are not only localized on the common organelles such as Golgi (TgDHH1, 5, 6, 9, 11, 12, 15 and 17), ER (TgDHH3, 8 and 16), PM (TgDHH4 and 13), but also the Apicomplexan-specific organelles such as IMC (TgDHH2 and 14) and rhoptries (TgDHH7) (Frénal et al., 2013).

In some contrast to the subcellular localization described above, studies carried out on transiently expressed in tobacco leaves of the 24 Arabidopsis PATs show that 9 of them are localized on PM, including AtPAT04-09, 12, 19 and 21. Therefore, it was proposed that PM is the main site for S-acylation in Arabidopsis plant (Batistic, 2012). This is different from mammalian PATs where most of them are localized in Golgi and therefore Golgi is thought to be the major S-acylation machinery (Ohno et al., 2006; Batistic, 2012). It is also interesting to note that many AtPATs that are localized on ER and PM are also associated with vesicles around them. For example, AtPAT3, 15, 17 and 18 are localized at ER as well as on the vesicles associated with them; AtPAT13, 20 and 22 at PM and also vesicles (Batistic, 2012). AtPAT10, 14, 16, 23 and 24 are mainly localized at Golgi, and the Golgi-localization of AtPAT10 and AtPAT14 were further confirmed in stably transformed Arabidopsis plants (Qi et al, 2013; Li et al, 2015). While the main residence of AtPAT01 and AtPAT02 are the endosomal compartments AtPAT10 and AtPAT11 were found on the tonoplast (Batistic, 2012; Qi et al, 2013; Zhou et al, 2013). It is noteworthy that a few Arabidopsis PATs have dual subcellular localizations, such as AtPAT10 (Golgi and tonoplast) and AtPAT13/20/22 (PM and vesicles) (Batistic, 2012; Qi et al., 2013). Similar observations were also made with some mammalian PATs (Valdez-Taubas and Pelham, 2005; Ohno et al., 2006). However, the significance of this dual-localization nature of PATs is currently unknown.

Little is known about how the PAT proteins achieve their respective localization in the cell. A recent study show that the lysine-based sorting signals KXX and KKXX are present in the mammalian DHH4 and DHH6, respectively, and it is these motifs that restrict their localization to the ER (Gorleku et al., 2011). It is also revealed that the C-

terminal 68 amino acids of the mammalian DHHC2 play an important role to define its subcellular localization to the ER and Golgi (Fukata et al., 2013). However, there is currently no information available on how plant PAT are targeted to individual membranes in the cell.

The Identified PAT/Substrate Pairs

As an enzyme PAT carries out its function mainly through substrate protein(s) it S-acylates. Therefore, to understand how PATs operate it is important to identify the target proteins they modify. However, to match an individual PAT and its S-acylated substrate proteins has proved to be a very difficult task so far. This is because: 1. The number of potential S-acylated proteins far exceed the number of their modifying PATs. For example, there are 7 PATs in yeast, however, ~50 S-acylated proteins were identified by a proteomic approach (Roth et al., 2006). Similarly, much more S-acylated proteins were isolated than the number of PATs present in mammals and Arabidopsis (Martin and Cravatt, 2009; Hemsley et al., 2013). Therefore, it seems most likely that at least some if not all PATs can S-acylate multiple substrate proteins, i.e., PATs do not have strict substrate specificity; 2. Many substrate proteins can be modified by more than one PATs. For instance, in yeast, the S-acylation of Vac8 is only partially reduced in the yeast PAT knock-out strain *pfa3*, thus it is most likely that Vac8 is S-acylated by Pfa3 as well as one another or other PATs (Smotrýs et al., 2005). Similarly, Ras2 S-acylation is only partially suppressed in the absence of Erf2 hence other PATs are also capable to S-acylate Ras2 (Roth et al., 2006; Gonzalez Montoro et al., 2011). Therefore, these PATs have specific yet overlapping substrate specificity. For some peripheral membrane proteins in mammalian cells, their S-acylation is devoid of specificity altogether (Rocks et al., 2010). However, reports show that some PATs do have their preferentially modified proteins. For example, Swf1 in yeast prefers to function on transmembrane proteins that have cysteines close to TMDs (Roth et al., 2006). In human, integrin $\alpha 4\beta 6$ is strictly modified by DHHC3 (Sharma et al., 2012); 3. No consensus sequences in S-acylated proteins have been found. Although many S-

acylated proteins have been identified and some of them are S-acylated by the same PAT, there are no consensus sequences characterised in these proteins (Gonzalez Montoro et al., 2011).

In yeast, each of the five PATs have been mapped to one or more substrate proteins. However, the total number of these individual substrate proteins are still far less than ~50 S-acylated proteins identified (Roth et al, 2006). For example, Akr1 S-acylates casein kinases Yck1, Yck2 and Yck3 (Roth et al., 2002). It also S-acylates sphingosine kinase Lcb4 because 60-80% of reduction in S-acylation of Lcb4 was found in *akr1* mutant yeast (Kihara et al., 2005). Other proteins that are also S-acylated by Akr1 are Meh1, Sna4 and the unknown function proteins such as Ypl199c, Ykl047w and Ypl236c (Roth et al., 2006). Therefore, Akr1 alone can S-acylate at least 9 substrate proteins in yeast. It was noted that Akr1 prefers hydrophilic proteins that tether to membranes solely through N- or C-terminal palmitoyl modifications (Roth et al., 2006). Erf2 is responsible for the S-acylation of Ras and other signalling proteins such as Rho2, Rho3, Gpa1, Gpa2 and Ste18, all of which are heterolipidated (Bartels et al., 1999; Lobo et al., 2002; Roth et al., 2006; Zhang et al., 2013). Swf1 tends to S-acylate proteins that have juxta-TMD mapping cysteines, such as SNAREs (Valdez-Taubas and Pelham, 2005), mannosyltransferases including Mnn1, Mnn10 and Mnn11 and prion induction protein Pin2 (Roth et al., 2006). Pfa4 is devoted to the palmitoylation of a group of Amino Acid Permeases (AAPs). AAPs is a family of plasma membrane transporters with 12 TMDs and a conserved C-terminal Phe-Trp-Cys palmitoylation site. Experiments in *Cryptococcus neoformans* showed that Pfa4 is also responsible for PM localization of Ras1 via palmitoylation (Merino et al., 2014). On the other hand, one substrate protein can be palmitoylated by multiple PATs. For example, S-acylation of Gpa2 is mediated by both Pfa5 and Erf2 (Roth et al., 2006; Zhang et al., 2013; Greaves and Chamberlain, 2011); Meh1 was S-acylated by Pfa3 and Akr1 (Greaves and Chamberlain, 2011); and an unknown protein Ygl108 was S-acylated equally by Erf2 and Pfa4 (Greaves and Chamberlain, 2011). However, so far the substrates of Akr2 has not been identified. Therefore, it is clear that both PATs and their substrate proteins are

highly redundant in yeast. A summary of PATs and their substrate proteins in yeast is shown in **Table 4**.

TABLE 4 | Substrates of Yeast PATs

PATs	Substrates	references
Akr1	Lcb4, Yck1, Yck2, Yck3, Meh1, Sna4, Ypl199c, Ykl047w, Ypl236c, Vac8	Roth et al., 2002; Babu et al., 2004; Kihara et al., 2005; Roth et al., 2006; Roth et al., 2011
Erf2 (shr5)	Ras1, Ras2, Rho2, Rho3, Gpa1, Gpa2, Ste18, Ycp4, Psr1, Yg1108	Bartels et al., 1999; Lobo et al., 2002; Ohno et al., 2006; Roth et al., 2006; Greaves and Chamberlain, 2011; Zhang et al., 2013;
Swf1	Many SNAREs, Mnn1, Mnn10, Mnn11, Pin2	Valdez-Taubas and Pelham, 2005; Roth et al., 2006
Pfa3	Vac8, Meh1	Hou et al., 2005; Smotrys et al., 2005; Roth et al., 2006
Pfa4	APPs, Lcb4, Ras1, Yg1108	Ohno et al., 2006; Roth et al., 2006; Greaves and Chamberlain, 2011; Nichols et al., 2015
Pfa5	Gpa2	Greaves and Chamberlain, 2011

Many substrate proteins of mammalian PATs have also been identified in recent years. These include GTP-binding proteins, cytoskeletal proteins, enzymes, neurotransmitter receptors and synaptic scaffolding proteins (**Table 5**). Similar to what is found in yeast, some proteins can be modified by more than one PAT and most PATs can modify multiple proteins such as DHHC2, 3, 5, 7, 8, 13, 15, 17 and 21 (**Table 5**). For instance, PSD-95, a protein that scaffolds receptors and signaling enzymes at the postsynapse (Topinka and Bredt, 1998) can be S-acylated by DHHC2, 3, 7, 8, 15 and 17 (Fukata et al., 2004; Fukata et al., 2006; Greaves and Chamberlain, 2011; Butland et al., 2014). SNAP-25, a t-SNARE protein that regulates neurotransmitter release, is the substrate of DHHC2, 3, 7, 8, 15 and 17 (Greaves et al., 2009). S-acylation of a tyrosine kinase Fyn is mediated by DHHC2, 3, 7, 15, 20 and 21 (Mill et al., 2009). All these mentioned PAT/substrates and other pairs are listed in **Table 5**. On the other hand, one PAT can palmitoylate multiple substrate proteins. For example, DHHC2

palmitoylates cytoskeleton-associated protein 4 (CKAP4) and AKAP79/150 (Chavda et al., 2014; Keith et al., 2012); DHHC3 does integrin $\alpha 6\beta 4$, Calmodulin-dependent protein kinase isoform 1 γ (CaMKI γ), NMDA receptor subunits 2A and 2B (NR2A/B) and DR4 (Sharma et al., 2012; Takemoto-Kimura et al., 2007; Hayashi, et al., 2009; Yeste-Velasco et al., 2015); the S-acylation of Grip1b, δ -catenin, Flotillin-2, somatostatin receptor 5 and Ankyrin-G is carried out by DHHC5 (Brigidi et al., 2015). In the same fashion many other proteins are also palmitoylated by specific PATs (**Table 5**). Importantly, some DHHC proteins have been indicated to be involved in certain diseases, such as DHHC8 in schizophrenia, DHHC9 and 15 in X-linked mental retardation, DHHC17 in Huntington's disease and many PATs are involved in different types of cancer including DHHC2, 3, 7, 9, 11, 14, 17, 20 and 21 (Chavda et al., 2014; Yeste-Velasco et al., 2015). However, for some of them, their specific substrate proteins have not been identified.

TABLE 5 | Mammalian PATs and their (regulated) target proteins

PAT	Targets	References
DHHC2	PSD-95, CKAP4, SNAP23/25, eNOS, Fyn, NDE1, NDEL1, CD9/151, ABCA1, AKAP79/150	Fukata et al., 2004; Huang et al., 2004; Fernandez-Hernando et al., 2006; Sharma et al., 2008; Zhang et al., 2008; Shmueli et al., 2009; Singaraja et al., 2009; Greaves et al., 2010; Chavda et al., 2014
DHHC3 (GODZ)	PSD-95, SNAP23/25/25b, G α , CSP, Integrin $\alpha 6\beta 4$, GABA γ 2, eNOS, GluR1/2, GAD65, STREX, Fyn, BACE1, NDE1, NDEL1, NCAM140, CaMKI γ , NR2A/B, DR4, PI4KII	Fukata et al., 2004; Hayashi et al., 2005; Fang et al., 2006; Fernandez-Hernando et al., 2006; Takemoto-Kimura et al., 2007; Greaves et al., 2008; Huang et al., 2009; Mill et al., 2009; Shmueli et al., 2009; Tsutsumi et al., 2009; Vetrivel et al., 2009; Greaves et al., 2010; Tian et al., 2010; Yeste-Velasco et al., 2015
DHHC4	BACE1	Vetrivel et al., 2009
DHHC5	Grip1b, δ -catenin, Flotillin-2, somatostatin receptor 5, Ankyrin-G, STREX	Tian et al., 2010; Kakkola et al., 2011; Li et al., 2012; Thomas et al., 2012; Brigidi et al., 2014; Brigidi et al., 2015
DHHC6	Chaperone calnexin	Lakkaraju et al., 2012
DHHC7	PSD-95, G α , CSP, Fyn, eNOS, SNAP25/23/25b, GABA γ 2, STREX,	Fukata et al., 2004; Fang et al., 2006; Fernandez-Hernando et al., 2006; Fukata et al., 2006; Greaves et al., 2008; McCormick et al., 2008; Ponimaskin et

	BACE1, NDE1, NDEL1, NCAM140, sortillin, PDE10A2, CD9, ER, PR, AR, PI4KII	al., 2008; Greaves et al., 2009; Shmueli et al., 2009; Tsutsumi et al., 2009; Vetrivel et al., 2009; Charych et al., 2010; Greaves et al., 2010; Tian et al., 2010; Ohno et al., 2012
DHHC8	eNOS, SNAP25, paralemmin-1, GAD65, PSD95, PSD93	Fernandez-Hernando et al., 2006; Mukai et al., 2008; Huang et al., 2009
DHHC9	H- and N-Ras, STREX	Swarthout et al., 2005; Tian et al., 2010
DHHC12	ABCA1	Singaraja et al., 2009; Chavda et al., 2014
DHHC13 (HIP14L)	MT1-MMP, HTT, GAD65	Huang et al., 2009; Saleem et al., 2010; Song et al., 2014
DHHC15	PSD95, GAP43, SNAP25b, CSP, GABA _A γ2, Fyn, BACE1, CD151, CI-MPR, sortillin	Fukata et al., 2004; Fang et al., 2006; Greaves et al., 2008; McCormick et al., 2008; Sharma et al., 2008; Mill et al., 2009; Vetrivel et al., 2009; Greaves et al., 2010
DHHC17 (HIP14)	PSD95, CLIP3, CSP, GAD65, GAP43, GLUR1/2, GPM6A, HTT, JNK3, Lck, SNAP25/23/25b, STREX, SYT1, SPRED1/3, Ras	Fukata et al., 2004; Greaves et al., 2008; Huang et al., 2009; Mill et al., 2009; Greaves et al., 2010; Tian et al., 2010; Ohno et al., 2012; Butland et al., 2014
DHHC18	H- and N-Ras, Lck	Fukata et al., 2004
DHHC19	R-Ras, PDE10A2	Charych et al., 2010; Baumgart et al., 2010
DHHC20	Fyn, BACE1, ABCA1	Mill et al., 2009; Singaraja et al., 2009; Vetrivel et al., 2009
DHHC21	PECAM1, SOD1, Lck, eNOS, Fyn, ABCA1, ER, PR, AR	Fernandez-Hernando et al., 2006; Takemoto-Kimura et al., 2007; Mill et al., 2009; Vetrivel et al., 2009; Antinone et al., 2013; Akimzhanov and Boehning, 2015; Yeste-Velasco et al., 2015

645

646 Very little information is available for PAT/substrate pairs in other organisms. The
647 only PAT/substrate pair characterized was in *Plasmodium falciparum* where PfDHHC1,
648 an apicomplexan-specific and inner membrane complex-localized PAT, has identical
649 expression pattern to two S-acylated proteins PfISP1 and PfISP3 (Wetzel et al., 2014).

In plant, the only putative PAT/substrates pairing identified is ATPAT10/AtCBL2, 3, 6. This was achieved by transient expression of AtCBL2, AtCBL3 and AtCBL6 in Arabidopsis protoplast, showing that the tonoplast localization of AtCBLs is lost in protoplast prepared from ATPAT10 loss-of-function mutant (Zhou et al., 2013).

Therefore, although many hundreds of S-acylated proteins, including putative ones isolated by large proteomic approaches were identified from different species at present, there are many more to come in the future due to the readily available proteomics facilities in large institutions. A framework for characterizing PAT/substrate selectivity is urgently required to set out to match individual PATs and their S-acylated substrate proteins in order to understand the mechanism of S-acylation in individual organism and in general.

DE-S-ACYLATION

Similar to phosphorylation and ubiquitination, S-acylation process is reversible, which makes it a very important lipid modification of proteins (Hemsley, 2009). S-acylation turnover by de-S-acylation, can be constitutive or stimulated (Smotrys and Linder, 2004). Ras proteins were the first proteins to be reported to have dynamic S-acylation with different H-Ras has different de-S-acylation rates (Baker et al., 2003). S-acylation/de-S-acylation of Fyn, a member of the Src kinase family, happens with a half-life of 1.5-2 hours (Wolven, et al., 1997; Zeidman et al., 2009). The de-S-acylation of G α subunits is stimulated by the activation of G-protein-coupled receptors (Mumby et al., 1994; Linder and Deschenes, 2007). De-S-acylation of PSD-95 is enhanced by neuronal activity (El-Husseini et al., 2002).

At present only four protein thioesterases have been identified to catalyze the de-S-acylation process, including acyl protein thioesterases 1 (APT1) and 2 (APT2), palmitoyl thioesterases 1 (PPT1) and 2 (PPT2) (Tomatis et al 2010; Hornemann, 2015).

These enzymes carry out the de-S-acylation step in which the palmitate or other long chain fatty acids are removed from the S-acylated proteins (Linder and Deschenes, 2007). APT1 was first found in rat liver as a lysophospholipase and its substrates include Ras, G α subunit, RGS4, SNAP-23 and eNOS (Toyoda et al., 1999; Akimzhanov and Boehning, 2015). APT2 was reported to de-S-acylate the growth-associated protein 43 (Tomatis et al., 2010). PPT1 is a soluble lipase that is localized in lysosomes and it is responsible for the degradation of S-acylated proteins (Linder and Deschenes, 2007; Chavda et al., 2014). The loss-of-function of PPT1 resulted in severe infantile neuronal ceroid lipofuscinosis (Vesa et al., 1995). PPT2 has very limited acyl protein thioesterase activity, which prefers de-S-acylating short-chain lipid substrate. Interestingly, study has shown that it is up-regulated in obesity (Burger et al., 2012; Fox et al., 2012).

It is surprising that only four thioesterases have been identified so far yet many hundreds of S-acylated proteins were isolated from different genomes. The explanations for this could be: 1) thioesterases are broad specificity enzymes, each of which can de-S-acylate a wide range of substrates; 2) not all S-acylated proteins undergo de-S-acylation; 3) of course, it could be because many more thioesterases have not been found (Chavda et al., 2014). There currently no protein thioesterases have been identified from plant.

MECHANISM OF PROTEIN S-ACYLATION

It is well recognized that DHHC proteins transfer acyl group via a two-step catalytic mechanism in which the enzyme first modifies itself with palmitate (or other long chain fatty acids) in a process termed autoacylation. The enzyme then transfers the acyl group from itself onto its substrate proteins. However, the number and location of the S-acylated cysteines of a given PAT in the autoacylated intermediate is unknown. It is well accepted that the cysteine in the DHHC motif is the auto-S-acylation site because mutation in this residue results in loss of auto-acylation of many characterized PATs from yeast (Montoro et al., 2011), mammals (Ohno et al., 2012) and Arabidopsis

(Hemsley et al., 2005; Qi et al., 2013). However, cysteines in other positions of PATs such as the CRD and other domains may also be autoacylated (Gottlieb et al., 2015). For instance, DHHC3 has 6 auto-S-acylation sites where 5 in the CRD, including Cys-132, Cys-133, Cys-146, Cys-157 and Cys-163, 1 in the N-terminal domain (Cys-24) (Gottlieb et al., 2015).

Techniques Used for Prediction and Confirmation of S-acylated Cysteines in Proteins

There is no consensus for sequences in the S-acylated proteins despite that many such proteins have been isolated through proteomics approach or individually confirmed via radioactive labelling or/and mutation studies. Nevertheless, it is noted that: 1) in some S-acylated soluble proteins the cysteine residues that are S-acylated are frequently surrounded by basic or hydrophobic amino acids, such as GAP-43 (Liu et al., 1993) at N-terminal motif, Yck2 (Roth et al., 2002) at C-terminal motif and SNAP-25b (Lane and Liu, 1997) at cysteine string motif (Smotrys and Linder, 2004); 2) in other S-acylated soluble proteins the Cys residue is located near the prenylated or myristoylated residues, resulting in the so-called dual lipidation. These proteins include the Arabidopsis α and γ subunits of heterotrimeric G protein (Adjobo-Hermans et al., 2006; Zeng et al., 2007), small GTPases (Roth et al., 2006; Zhang et al., 2013; Deschenes et al., 1990; Bartels et al., 1999); 3) for transmembrane proteins, the Cys residues are often situated in the cytoplasmic regions of membrane-spanning regions (Roth et al., 2006; Ohno et al., 2012). For instance, the S-acylation of C261-263 triplet in death receptor 4 (DR4) (Rossin et al., 2009) and C474 in β -secretase BACE1 (Motoki et al., 2012) promotes their association with lipid raft.

Based on the above information several software packages have been developed to predict the S-acylated cysteines, such as a clustering and scoring strategy known as CSS-Palm (Zhou et al., 2006), which has been updated to the latest version CSS-Palm 4.0 (freely available at <http://csspalm.biocuckoo.org/>), incremental feature selection (IFS)-Palm (Hu et al., 2011), weight, amino acid composition and position specific

scoring (WAP)-Palm (Shi et al., 2013) and PalmPred (Kumari et al., 2014). All of them are on-line so that one can input the protein sequence of interest to predict the possibility of its S-acylation and where the Cys residues are located within the sequence. The prediction data from these platforms can then be confirmed experimentally. These techniques include:

1. PAT inhibitors. The palmitate analog, 2-bromopalmitate (2-BP) is the most commonly used inhibitor of S-acylation, which inhibits palmitoylation in cells and PAT activity of DHHC proteins *in vitro* (Webb et al., 2000; Fukata et al., 2004; Jennings et al., 2009). However, it lacks specificity and can also inhibit myristoylation and reduce de-acylation through inhibiting activities of acyl-protein thioesterases (Webb et al., 2000; Pedro et al., 2013). Tunicamycin and cerulenin are also used to inhibit S-acylation, but similar effect was found as 2-BP (Patterson and Skene, 1995; Lawrence et al., 1999). Recently, a compound, 2-(2-hydroxy-5-nitro-benzylidene)-benzo[*b*]thiophen-3-one, was shown to have more specificity, but it does not have selectivity for specific PAT, i.e., it inhibits activities of all PATs (Jennings et al., 2009), which means it still cannot be used to study the function of individual PAT. Therefore, it is clear that results obtained from these inhibitors should be further validated by mutational or biochemical analysis.

2. Mutational analysis to change the potential S-acylated Cysteine to alanine or serine. Both alanine and serine were frequently used to replace the cysteine to achieve similar results (Hemsley et al., 2005; Qi et al., 2013; Li et al., 2015). Cysteine and serine have very similar structure, when the cysteine is mutated to serine it can maintain the size and the properties of the putative S-acylated protein, in this case, serine is a better substitution for cysteine. However, compare to alanine, serine is more hydrophilic than cysteine and might also cause unwanted side chain effects (Nagano et al., 1999). In this specific study, both alanine and serine as the substitutions for cysteine are accepted so far. This is followed by comparing the effect on the differences of functions or the subcellular localizations to native protein. If a difference was found the proteins were

most likely S-acylated at the cysteine residues that were mutated.

3. Biochemical assays to analyze the attachment of fatty acids of the individual proteins. This includes: 1) traditionally feed with tritiated fatty acids followed by exposures to X-ray film (Lavy et al., 2002); 2) azido-alkyne CLICK-chemistry (Martin & Cravatt, 2009); 3) Acyl-exchange, or Biotin-switch assay (Wan et al., 2007; Hemsley et al., 2008); and 4) direct resin immobilisation (Forrester et al., 2010).

4. Direct detection of the S-acyl group by gas chromatography–mass spectrometry (GC-MS) analysis. The identification of lipid groups attached to proteins can help to understand the biophysical properties of the protein. This method has been successfully used to demonstrate S-acylation of CBL1 and CBL2, which are attached by palmitate and/or stearate (Batistic et al., 2008; Batistic et al., 2012).

SPECIFICITIES OF PAT-SUBSTRATE INTERACTION

Although it is generally accepted that PATs are lacking specificity towards their substrate proteins and vice versa studies in yeast showed that some PATs do exhibit preference to some substrate protein(s) compared than others. For instance, Akr1 prefers to S-acylate soluble proteins at their N- or C-terminus, such as Ypl236c S-acylated at N-terminal cysteine and Yck1 S-acylated at C-terminal cysteine. Erf2 and Pfa5 show preference for pre-lipidated substrates, such as prenylated Ras1 and Ras2, myristoylated Gpa1 and Gpa2. Swf1 and Pfa4, on the other hand, prefer single and multiple transmembrane proteins such as SNAREs and AAPs (Roth et al., 2006; Ohno et al., 2012). Similar conclusions were made from studies of mammalian DHHC proteins where it was found that DHHC3, 7, 8 and 14-17 had high activities towards soluble proteins, while DHHC2, 20 and 21 were highly active to integral membrane proteins (Ohno et al., 2012). However, it was also noted that most DHHC proteins in both yeast and mammals had overlap activity to modify pre-lipidated substrates, such

as 4 yeast PATs and 16 mammalian PATs can all S-acylate the myristoylated Gpa2 (Ohno et al., 2012).

The question here is how PATs and their substrates recognize each other? To address this, studies were carried out on some PATs and their S-acylated proteins in mammalian system. It was reported that the AR domain of the two ankyrin repeat containing PATs DHHC13 and 17 in mammals can act as substrate-recruiting signal and recognizes the [VIAP][VIT]XXQP motif that is shared between some S-acylated proteins including SNAP25/23, CSP, HTT and CLIP3 (Lemonidis et al., 2015). Fusing the AR domain of DHHC17 to the N terminus of DHHC3 that lacks an AR domain, can make DHHC3 a PAT for HTT which also supports the notion that the AR domain contributes to the substrate specificity of DHHC17 for HTT (Huang et al., 2009). DHHC3 and 7 interact with GABA_Aγ2 through a 14-amino acid cysteine rich domain (Fang et al., 2006). DHHC7 has two splicing isoforms, the longer one has additional 111bp compared to the shorter one, which might possess its tissue-specific function since it expresses specifically in placenta, lung, liver, thymus and small intestine (Ohno et al., 2006). The recognition and S-acylation of PSD95 by DHHC17 depend on the N-terminal 13 amino acids of PSD-95 (Huang et al., 2009). The cysteine rich ‘CCPCC’ motif of PI4KII is required for its S-acylation by DHHC3 and DHHC7 (Lu et al., 2012). Subtle changes in the S-acylation domains of proteins can alter their PAT specificity, which were proved from SNAP23/25 (Greaves et al., 2010). For instance, a SNAP25 mutant which lacks a proline located 25 residues downstream of the S-acylated domain can only be modified by DHHC3 but not DHHC17 (Greaves et al., 2009). Therefore, specific domains in PATs and their substrate proteins are required for recognition and S-acylation to occur.

Subcellular localizations of PATs can have a profound effect on the type of proteins it can S-acylate (Greaves and Chamberlain, 2011). A transmembrane protein might only have access to be S-acylated by the PATs localized on the same membrane. For example, the PM-localized Gpa1 is S-acylated by the PM-localized Pfa5 in yeast (Ohno et al.,

2006); tonoplast-localization of AtCBL2 and AtCBL3 is via S-acylation carried out by AtPAT10 which is also localized on tonoplast in Arabidopsis (Zhou et al., 2013).

For S-acylation of a protein to occur its prior membrane attachment via TMD, another lipid modification or protein-protein interaction is often acquired (Hemsley, 2015). For instance, TEM8 localizes at PM with one TMD, S-acylation of which negatively regulate its raft association (Abrami et al., 2006). Some proteins require another lipid modification such as myristoylation to target to certain membrane first before the S-acylation can take place. AtCBL1 is one of these proteins where myristoylation targets it on the ER, then the unknown ER-localized PAT S-acylates the myristoylated AtCBL1. This dual-lipidated AtCBL1 can subsequently be trafficked to the PM (Batistič et al., 2008). It was reported that the N-terminal 12-amino acid peptide of AtCBLs is sufficient to mediate the dual lipid modification and target to PM or tonoplast (Batistic et al., 2008). Therefore, the localization of a specific PAT for a given S-acylated substrate protein relies on where this protein is localized after the first lipid modification.

Important Molecules That Are Involved In S-Acylation

Special molecules have either positive or negative effect on S-acylation of certain proteins. These molecules could be another protein, hormone, ions or protein inhibitors. For instance, although most DHHC-PATs in mammals can catalyse S-acylation independently DHHC9 needs a Golgi-localized protein GCP16 to specially palmitoylate H- and N-Ras (Swarthout et al., 2005). Ykt6, which possibly works as a co-factor of Pfa3 enhanced the S-acylation and vacuole localization of Vac8 (Dietrich et al., 2004; Hou et al., 2005; Meiringer and Ungermann, 2006). S-acylation and localization of Ras protein is catalysed by the Erf2p-Erf4p complex in yeast (Zhao et al., 2002). Zinc ion is tightly bound to the cysteine rich domain of the DHHC3, which is essential for its structural integrity and PAT activity (Gottlieb et al., 2015). Selenoprotein K (SelK), an 11-kDa endoplasmic reticulum protein of unknown function (Shchedrina et al, 2011) is required for the S-acylation of both IP3R (inositol-1, 4, 5-

triphosphate receptor) and CD36 (Fredericks and Hoffmann, 2015). S-acylation of PI4KII is cholesterol-dependent (Lu et al., 2012). Some compounds might have negative effect to specific S-acylation, such as curcumin can prevent S-acylation of integrin $\beta 4$ by DHHC3 in breast cancer cells (Coleman et al., 2015). Understanding the involvement of these molecules in S-acylation could provide important information in designing and developing new drugs to target the disease and cancer-related S-acylation machinery.

CONCLUSION AND FUTURE PERSPECTIVES

Ever since the discovery of the first S-acyltransferase, Akr1 from yeast in 2002, which lead to the realization of protein S-acylation being an enzymatic process rather than a simultaneous addition of a long chain fatty acid to proteins, research on S-acylation of proteins has accelerated in a remarkable speed in the past decade. Yeast, as a simple unicellular model eukaryote, has been the first choice for researchers to study S-acylation. The knowledge learnt from the yeast system has then been applied in guiding similar studies in other organisms. As such the important roles of protein S-acylation in growth and development, especially in different human diseases, such as cancers, have hence attracted much attention and become a hot area of research in recent years.

Although progress has been made toward understanding various aspects of protein palmitoylation the corresponding research in plants is trailing behind that in yeast and mammals. Judging from the wide arrays of S-acylated proteins identified recently from Arabidopsis and Poplar by proteomic studies it is clear that S-acylation plays variable and important roles in plant growth, development and environmental adaption (Hemsley et al, 2013; Srivastava et al., 2016). The knowledge learnt and methodologies developed from yeast and mammals will no doubt provide important clues and necessary tools for us to conduct more efficient research on S-acylation in plants in the coming years. Specifically, 1. the roles of the remaining 21 AtPATs in Arabidopsis will need to be characterized. PATs from other plant species, such as poplar, especially with

its S-acylated proteins being isolated recently, will also need to be characterized to see if plant PATs share functional similarity, this will further validate the data obtained from Arabidopsis PATs so far; 2. To match individual PATs with their S-acylated substrate proteins in Arabidopsis and poplar. At present, the only plant PAT with tentative mapped substrate proteins is the Arabidopsis AtPAT10 where it was found that the tonoplast localization of transiently expressed CBL2,3,6 were lost in the protoplast prepared from leaf cells of AtPAT10 loss-of-function mutant plant (Zhou et al, 2013). This indicates that AtPAT10 functions in calcium signaling and salt stress through the actions of these CBLs. Similar approaches could be used to map other PAT/substrate(s) pairs in Arabidopsis and poplar. This will provide further insights to substrate specificity of PATs and molecular mechanisms how PATs function in plants; 3. De-palmitoylation enzymes. S-acylation of proteins is a reversible process where S-acylation is catalyzed by PATs and De-palmitoylation is by acyl protein thioesterases. While 4 such enzymes have been identified and characterized from mammals none from plant. Therefore, research in this area is paramount.

REFERENCES

- Abrami, L., Kunz, B., Iacovache, I., and Van Der Goot, F.G. (2008). Palmitoylation and ubiquitination regulate exit of the Wnt signaling protein LRP6 from the endoplasmic reticulum. *Proceedings of the National Academy of Sciences* 105, 5384-5389.
- Abrami, L., Leppla, S.H., and Van Der Goot, F.G. (2006). Receptor palmitoylation and ubiquitination regulate anthrax toxin endocytosis. *The Journal of cell biology* 172, 309-320.
- Adams, M., Harrington, B., He, Y., Davies, C., Wallace, S., Chetty, N., Crandon, A., Oliveira, N., Shannon, C., and Coward, J. (2015). EGF inhibits constitutive internalization and palmitoylation-dependent degradation of membrane-spanning procancer CDCP1 promoting its availability on the cell surface. *Oncogene* 34, 1375-1383.
- Adjobo-Hermans, M.J., Goedhart, J., and Gadella, T.W. (2006). Plant G protein heterotrimers require dual lipidation motifs of G α and G γ and do not dissociate upon activation. *Journal of cell science* 119, 5087-5097.
- Akimzhanov, A.M., and Boehning, D. (2015). Rapid and transient palmitoylation of the tyrosine kinase Lck mediates Fas signaling. *Proceedings of the National Academy of Sciences* 112, 11876-11880.

- Al Abdallah, Q., and Fortwendel, J.R. (2015). Exploration of *Aspergillus fumigatus* Ras pathways for novel antifungal drug targets. *Frontiers in microbiology* 6.
- Alassimone, J., Fujita, S., Doblas, V.G., Van Dop, M., Barberon, M., Kalmbach, L., Vermeer, J.E., Rojas-Murcia, N., Santuari, L., and Hardtke, C.S. (2016). Polarly localized kinase SGN1 is required for Casparian strip integrity and positioning. *Nature Plants* 2, 16113.
- Antinone, S.E., Ghadge, G.D., Lam, T.T., Wang, L., Roos, R.P., and Green, W.N. (2013). Palmitoylation of superoxide dismutase 1 (SOD1) is increased for familial amyotrophic lateral sclerosis-linked SOD1 mutants. *Journal of Biological Chemistry* 288, 21606-21617.
- Babu, P., Deschenes, R.J., and Robinson, L.C. (2004). Akr1p-dependent palmitoylation of Yck2p yeast casein kinase 1 is necessary and sufficient for plasma membrane targeting. *Journal of Biological Chemistry* 279, 27138-27147.
- Baekkeskov, S., and Kanaani, J. (2009). Palmitoylation cycles and regulation of protein function (Review). *Molecular membrane biology* 26, 42-54.
- Baker, T.L., Zheng, H., Walker, J., Coloff, J.L., and Buss, J.E. (2003). Distinct rates of palmitate turnover on membrane-bound cellular and oncogenic H-ras. *Journal of Biological Chemistry* 278, 19292-19300.
- Bannan, B.A., Van Etten, J., Kohler, J.A., Tsoi, Y., Hansen, N.M., Sigmon, S., Fowler, E., Buff, H., Williams, T.S., and Ault, J.G. (2008). The *Drosophila* protein palmitoylome: characterizing palmitoyl-thioesterases and DHHC palmitoyl-transferases. *Fly* 2, 198-214.
- Bartels, D.J., Mitchell, D.A., Dong, X., and Deschenes, R.J. (1999). Erf2, a novel gene product that affects the localization and palmitoylation of Ras2 in *Saccharomyces cerevisiae*. *Molecular and cellular biology* 19, 6775-6787.
- Batistič, O. (2012). Genomics and Localization of the Arabidopsis DHHC-Cysteine-Rich Domain S-Acyltransferase Protein Family. *Plant Physiology* 160, 1597-1612.
- Batistič, O., Rehers, M., Akerman, A., Schlücking, K., Steinhorst, L., Yalovsky, S., and Kudla, J. (2012). S-acylation-dependent association of the calcium sensor CBL2 with the vacuolar membrane is essential for proper abscisic acid responses. *Cell research* 22, 1155-1168.
- Batistič, O., Sorek, N., Schültke, S., Yalovsky, S., and Kudla, J. (2008). Dual fatty acyl modification determines the localization and plasma membrane targeting of CBL/CIPK Ca²⁺ signaling complexes in Arabidopsis. *The Plant Cell* 20, 1346-1362.
- Baumgart, F., Corral-Escariz, M., Pérez-Gil, J., and Rodríguez-Crespo, I. (2010). Palmitoylation of R-Ras by human DHHC19, a palmitoyl transferase with a CaaX box. *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1798, 592-604.
- Baumgart, F., Corral-Escariz, M., Pérez-Gil, J., and Rodríguez-Crespo, I. (2010). Palmitoylation of R-Ras by human DHHC19, a palmitoyl transferase with a CaaX box. *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1798, 592-604.
- Beck, J.R., Rodriguez-Fernandez, I.A., De Leon, J.C., Huynh, M.-H., Carruthers, V.B., Morrisette, N.S., and Bradley, P.J. (2010). A novel family of *Toxoplasma* IMC proteins displays a hierarchical organization and functions in coordinating parasite division. *PLoS Pathog* 6, e1001094.

946 Benjannet, S., Elagoz, A., Wickham, L., Mamarbachi, M., Munzer, J.S., Basak, A., Lazure, C.,
 947 Cromlish, J.A., Sisodia, S., and Checler, F. (2001). Post-translational Processing of β -
 948 Secretase (β -Amyloid-converting Enzyme) and Its Ectodomain Shedding THE PRO-
 949 AND TRANSMEMBRANE/CYTOSOLIC DOMAINS AFFECT ITS CELLULAR
 950 ACTIVITY AND AMYLOID- β PRODUCTION. *Journal of Biological Chemistry* 276,
 951 10879-10887.

952 Bijlmakers, M.-J. (2009). Protein acylation and localization in T cell signaling (Review).
 953 *Molecular membrane biology* 26, 93-103.

954 Bishop, R.E. (2005). The lipid A palmitoyltransferase PagP: molecular mechanisms and role in
 955 bacterial pathogenesis. *Molecular microbiology* 57, 900-912.

956 Bishop, R.E., Kim, S.-H., and El Zoeiby, A. (2005). Role of lipid A palmitoylation in bacterial
 957 pathogenesis. *Journal of endotoxin research* 11, 174-180.

958 Bizzozero, O.A., Bixler, H.A., and Pastuszyn, A. (2001). Structural determinants influencing
 959 the reaction of cysteine-containing peptides with palmitoyl-coenzyme A and other
 960 thioesters. *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular*
 961 *Enzymology* 1545, 278-288.

962 Blaskovic, S., Blanc, M., and Goot, F.G. (2013). What does S-palmitoylation do to membrane
 963 proteins? *FEBS Journal* 280, 2766-2774.

964 Boyle, P.C., Schwizer, S., Hind, S.R., Kraus, C.M., Diaz, S.T., He, B., Martin, G.B. (2016).
 965 Detecting N-myristoylation and S-acylation of host and pathogen proteins in plants
 966 using click chemistry. *Plant Methods* 12: 38.

967 Brett, K., Kordyukova, L.V., Serebryakova, M.V., Mintaev, R.R., Alexeevski, A.V., and Veit,
 968 M. (2014). Site-specific S-Acylation of Influenza Virus Hemagglutinin THE
 969 LOCATION OF THE ACYLATION SITE RELATIVE TO THE MEMBRANE
 970 BORDER IS THE DECISIVE FACTOR FOR ATTACHMENT OF STEARATE.
 971 *Journal of Biological Chemistry* 289, 34978-34989.

972 Brigidi, G.S., Santyr, B., Shimell, J., Jovellar, B., and Bamji, S.X. (2015). Activity-regulated
 973 trafficking of the palmitoyl-acyl transferase DHHC5. *Nature communications* 6.

974 Brigidi, G.S., Sun, Y., Beccano-Kelly, D., Pitman, K., Mobasser, M., Borgland, S.L.,
 975 Milnerwood, A.J., and Bamji, S.X. (2014). Palmitoylation of Δ -catenin by
 976 DHHC5 mediates activity-induced synapse plasticity. *Nature neuroscience* 17, 522-
 977 532.

978 Buglino, J.A., and Resh, M.D. (2012). Palmitoylation of Hedgehog proteins. *Vitamins and*
 979 *hormones* 88, 229.

980 Bürger, M., Zimmermann, T.J., Kondoh, Y., Stege, P., Watanabe, N., Osada, H., Waldmann,
 981 H., and Vetter, I.R. (2012). Crystal structure of the predicted phospholipase LYPLAL1
 982 reveals unexpected functional plasticity despite close relationship to acyl protein
 983 thioesterases. *Journal of lipid research* 53, 43-50.

984 Butland, S.L., Sanders, S.S., Schmidt, M.E., Riechers, S.-P., Lin, D.T., Martin, D.D., Vaid, K.,
 985 Graham, R.K., Singaraja, R.R., and Wanker, E.E. (2014). The palmitoyl acyltransferase
 986 HIP14 shares a high proportion of interactors with huntingtin: implications for a role
 987 in the pathogenesis of Huntington's disease. *Human molecular genetics* 23, 4142-4160.

988 Casey, P.J., and Seabra, M.C. (1996). Protein prenyltransferases. *Journal of Biological*
 989 *Chemistry* 271, 5289-5292.

- Cavalier-Smith, T. (1993). Kingdom protozoa and its 18 phyla. *Microbiological reviews* 57, 953-994.
- Chakrabandhu, K., Hérincs, Z., Huault, S., Dost, B., Peng, L., Conchonaud, F., Marguet, D., He, H.T., and Hueber, A.O. (2007). Palmitoylation is required for efficient Fas cell death signaling. *The EMBO journal* 26, 209-220.
- Chamoun, Z., Mann, R.K., Nellen, D., Von Kessler, D.P., Bellotto, M., Beachy, P.A., and Basler, K. (2001). Skinny hedgehog, an acyltransferase required for palmitoylation and activity of the hedgehog signal. *Science* 293, 2080-2084.
- Charollais, J., and Van Der Goot, F.G. (2009). Palmitoylation of membrane proteins (Review). *Molecular membrane biology* 26, 55-66.
- Charych, E.I., Jiang, L.-X., Lo, F., Sullivan, K., and Brandon, N.J. (2010). Interplay of palmitoylation and phosphorylation in the trafficking and localization of phosphodiesterase 10A: implications for the treatment of schizophrenia. *The Journal of neuroscience* 30, 9027-9037.
- Chavda, B., Arnott, J.A., and Planey, S.L. (2014). Targeting protein palmitoylation: selective inhibitors and implications in disease. *Expert opinion on drug discovery* 9, 1005-1019.
- Chen, Z.H., Lam, H.C., Jin, Y., Kim, H.P., Cao, J., Lee, S.J., Ifedigbo, E., Parameswaran, H., Ryter, S.W., and Choi, A.M. (2010). Autophagy protein microtubule-associated protein 1 light chain-3B (LC3B) activates extrinsic apoptosis during cigarette smoke-induced emphysema. *Proceedings of the National Academy of Sciences* 107, 18880-18885.
- Cocca, S.M. (2014). *Phylogenetic analysis, modeling and experimental studies of the Saccharomyces cerevisiae palmitoylated protein kinase gene, ENV7*. California State University, Long Beach.
- Cohen, E. (1993). Chitin synthesis and degradation as targets for pesticide action. *Archives of insect biochemistry and physiology* 22, 245-261.
- Coleman, D.T., Soung, Y.H., Surh, Y.-J., Cardelli, J.A., and Chung, J. (2015). Curcumin prevents palmitoylation of integrin $\beta 4$ in breast cancer cells. *PloS one* 10, e0125399.
- Couve, A., Protopopov, V., and Gerst, J.E. (1995). Yeast synaptobrevin homologs are modified posttranslationally by the addition of palmitate. *Proceedings of the National Academy of Sciences* 92, 5987-5991.
- Deschenes, R., Resh, M., and Broach, J. (1990). Acylation and prenylation of proteins. *Current opinion in cell biology* 2, 1108-1113.
- Dietrich, L.E., and Ungermann, C. (2004). On the mechanism of protein palmitoylation. *EMBO reports* 5, 1053-1057.
- Dietrich, L.E., Gurezka, R., Veit, M., and Ungermann, C. (2004). The SNARE Ykt6 mediates protein palmitoylation during an early stage of homotypic vacuole fusion. *The EMBO journal* 23, 45-53.
- Dowen, R.H., Engel, J.L., Shao, F., Ecker, J.R., and Dixon, J.E. (2009). A family of bacterial cysteine protease type III effectors utilizes acylation-dependent and-independent strategies to localize to plasma membranes. *Journal of Biological Chemistry* 284, 15867-15879.
- Edmonds, M.J., and Morgan, A. (2014). A systematic analysis of protein palmitoylation in *Caenorhabditis elegans*. *BMC genomics* 15, 841.

- El-Husseini, A.E.D., Schnell, E., Dakoji, S., Sweeney, N., Zhou, Q., Prange, O., Gauthier-Campbell, C., Aguilera-Moreno, A., Nicoll, R.A., and Brecht, D.S. (2002). Synaptic strength regulated by palmitate cycling on PSD-95. *Cell* 108, 849-863.
- Fang, C., Deng, L., Keller, C.A., Fukata, M., Fukata, Y., Chen, G., and Lüscher, B. (2006). GODZ-mediated palmitoylation of GABAA receptors is required for normal assembly and function of GABAergic inhibitory synapses. *The Journal of neuroscience* 26, 12758-12768.
- Fernández-Hernando, C., Fukata, M., Bernatchez, P.N., Fukata, Y., Lin, M.I., Brecht, D.S., and Sessa, W.C. (2006). Identification of Golgi-localized acyl transferases that palmitoylate and regulate endothelial nitric oxide synthase. *The Journal of cell biology* 174, 369-377.
- Flannery, A.R., Czibener, C., and Andrews, N.W. (2010). Palmitoylation-dependent association with CD63 targets the Ca²⁺ sensor synaptotagmin VII to lysosomes. *The Journal of cell biology* 191, 599-613.
- Forrester, M.T., Hess, D.T., Thompson, J.W., Hultman, R., Moseley, M.A., Stamler, J.S., and Casey, P.J. (2011). Site-specific analysis of protein S-acylation by resin-assisted capture. *Journal of lipid research* 52, 393-398.
- Fox, C.S., Liu, Y., White, C.C., Feitosa, M., Smith, A.V., Heard-Costa, N., Lohman, K., Johnson, A.D., Foster, M.C., and Greenawalt, D.M. (2012). Genome-wide association for abdominal subcutaneous and visceral adipose reveals a novel locus for visceral fat in women. *PLoS Genet* 8, e1002695.
- Fredericks, G.J., and Hoffmann, P.R. (2015). Selenoprotein K and protein palmitoylation. *Antioxidants & redox signaling* 23, 854-862.
- Fréchal, K., Kemp, L.E., and Soldati-Favre, D. (2014). Emerging roles for protein S-palmitoylation in Toxoplasma biology. *International journal for parasitology* 44, 121-131.
- Fréchal, K., Polonais, V., Marq, J.-B., Stratmann, R., Limenitakis, J., and Soldati-Favre, D. (2010). Functional dissection of the apicomplexan glideosome molecular architecture. *Cell host & microbe* 8, 343-357.
- Fréchal, K., Tay, C.L., Mueller, C., Bushell, E.S., Jia, Y., Graindorge, A., Billker, O., Rayner, J.C., and Soldati-Favre, D. (2013). Global Analysis of Apicomplexan Protein S-Acyl Transferases Reveals an Enzyme Essential for Invasion. *Traffic* 14, 895-911.
- Fröhlich, M., Dejanovic, B., Kashkar, H., Schwarz, G., and Nussberger, S. (2014). S-palmitoylation represents a novel mechanism regulating the mitochondrial targeting of BAX and initiation of apoptosis. *Cell death & disease* 5, e1057.
- Fukasawa, M., Varlamov, O., Eng, W.S., Söllner, T.H., and Rothman, J.E. (2004). Localization and activity of the SNARE Ykt6 determined by its regulatory domain and palmitoylation. *Proceedings of the National Academy of Sciences of the United States of America* 101, 4815-4820.
- Fukata, M., Fukata, Y., Adesnik, H., Nicoll, R.A., and Brecht, D.S. (2004). Identification of PSD-95 palmitoylating enzymes. *Neuron* 44, 987-996.
- Fukata, Y., and Fukata, M. (2010). Protein palmitoylation in neuronal development and synaptic plasticity. *Nature Reviews Neuroscience* 11, 161-175.

1076 Fukata, Y., Dimitrov, A., Boncompain, G., Vielemeyer, O., Perez, F., and Fukata, M. (2013).
1077 Local palmitoylation cycles define activity-regulated postsynaptic subdomains. *The*
1078 *Journal of cell biology* 202, 145-161.

1079 Fukata, Y., Iwanaga, T., and Fukata, M. (2006). Systematic screening for palmitoyl transferase
1080 activity of the DHHC protein family in mammalian cells. *Methods* 40, 177-182.

1081 Furne, C., Corset, V., Hérincs, Z., Cahuzac, N., Hueber, A.-O., and Mehlen, P. (2006). The
1082 dependence receptor DCC requires lipid raft localization for cell death signaling.
1083 *Proceedings of the National Academy of Sciences of the United States of America* 103,
1084 4128-4133.

1085 Gagne, J.M., and Clark, S.E. (2010). The Arabidopsis stem cell factor POLTERGEIST is
1086 membrane localized and phospholipid stimulated. *The Plant Cell* 22, 729-743.

1087 Galli, L.M., Barnes, T.L., Secrest, S.S., Kadowaki, T., and Burrus, L.W. (2007). Porcupine-
1088 mediated lipid-modification regulates the activity and distribution of Wnt proteins in
1089 the chick neural tube. *Development* 134, 3339-3348.

1090 Ganesan, L., and Levental, I. (2015). Pharmacological Inhibition of Protein Lipidation. *The*
1091 *Journal of membrane biology* 248, 929-941.

1092 Gargantini, P.R., Gonzalez-Rizzo, S., Chinchilla, D., Raices, M., Giammaria, V., Ulloa, R.M.,
1093 Frugier, F., and Crespi, M.D. (2006). A CDPK isoform participates in the regulation of
1094 nodule number in *Medicago truncatula*. *The Plant Journal* 48, 843-856.

1095 Gaughran, J.P., Lai, M.H., Kirsch, D.R., and Silverman, S.J. (1994). Nikkomycin Z is a specific
1096 inhibitor of *Saccharomyces cerevisiae* chitin synthase isozyme Chs3 in vitro and in
1097 vivo. *Journal of Bacteriology* 176, 5857-5860.

1098 Gleason, E.J., Lindsey, W.C., Kroft, T.L., Singson, A.W., and L'hernault, S.W. (2006). spe-10
1099 encodes a DHHC-CRD zinc-finger membrane protein required for endoplasmic
1100 reticulum/Golgi membrane morphogenesis during *Caenorhabditis elegans*
1101 spermatogenesis. *Genetics* 172, 145-158.

1102 Gonzalez, M.A., Quiroga, R., Maccioni, H., and Valdez, T.J. (2009). A novel motif at the C-
1103 terminus of palmitoyltransferases is essential for Swf1 and Pfa3 function in vivo.
1104 *Biochem. J* 419, 301-308.

1105 Gorleku, O.A., Barns, A.-M., Prescott, G.R., Greaves, J., and Chamberlain, L.H. (2011).
1106 Endoplasmic reticulum localization of DHHC palmitoyltransferases mediated by
1107 lysine-based sorting signals. *Journal of Biological Chemistry* 286, 39573-39584.

1108 Gottlieb, C.D., Zhang, S., and Linder, M.E. (2015). The Cysteine-rich Domain of the DHHC3
1109 Palmitoyltransferase Is Palmitoylated and Contains Tightly Bound Zinc. *Journal of*
1110 *Biological Chemistry* 290, 29259-29269.

1111 Grassie, M.A., McCallum, J., Guzzi, F., Magee, A., Milligan, G., and Parenti, M. (1994). The
1112 palmitoylation status of the G-protein Go1 α regulates its activity of interaction with
1113 the plasma membrane. *Biochemical Journal* 302, 913-920.

1114 Greaves, J., and Chamberlain, L.H. (2007). Palmitoylation-dependent protein sorting. *The*
1115 *Journal of cell biology* 176, 249-254.

1116 Greaves, J., and Chamberlain, L.H. (2010). S-acylation by the DHHC protein family.
1117 *Biochemical Society Transactions* 38, 522-524.

1118 Greaves, J., and Chamberlain, L.H. (2011). DHHC palmitoyl transferases: substrate
1119 interactions and (patho) physiology. *Trends in biochemical sciences* 36, 245-253.

1120 Greaves, J., Gorleku, O.A., Salaun, C., and Chamberlain, L.H. (2010). Palmitoylation of the
 1121 SNAP25 Protein Family SPECIFICITY AND REGULATION BY DHHC
 1122 PALMITOYL TRANSFERASES. *Journal of Biological Chemistry* 285, 24629-24638.

1123 Greaves, J., Prescott, G.R., Fukata, Y., Fukata, M., Salaun, C., and Chamberlain, L.H. (2009).
 1124 The hydrophobic cysteine-rich domain of SNAP25 couples with downstream residues
 1125 to mediate membrane interactions and recognition by DHHC palmitoyl transferases.
 1126 *Molecular biology of the cell* 20, 1845-1854.

1127 Greaves, J., Salaun, C., Fukata, Y., Fukata, M., and Chamberlain, L.H. (2008). Palmitoylation
 1128 and membrane interactions of the neuroprotective chaperone cysteine-string protein.
 1129 *Journal of Biological Chemistry* 283, 25014-25026.

1130 Guardiola-Serrano, F., Rossin, A., Cahuzac, N., Lückcrath, K., Melzer, I., Mailfert, S., Marguet,
 1131 D., Zörnig, M., and Hueber, A. (2010). Palmitoylation of human FasL modulates its
 1132 cell death-inducing function. *Cell death & disease* 1, e88.

1133 Harada, T., Matsuzaki, O., Hayashi, H., Sugano, S., Matsuda, A., and Nishida, E. (2003).
 1134 AKRL1 and AKRL2 activate the JNK pathway. *Genes to Cells* 8, 493-500.

1135 Harashima, T., and Heitman, J. (2005). Gα subunit Gpa2 recruits kelch repeat subunits that
 1136 inhibit receptor-G protein coupling during cAMP-induced dimorphic transitions in
 1137 *Saccharomyces cerevisiae*. *Molecular biology of the cell* 16, 4557-4571.

1138 Hayashi, T., Rumbaugh, G., and Haganir, R.L. (2005). Differential regulation of AMPA
 1139 receptor subunit trafficking by palmitoylation of two distinct sites. *Neuron* 47, 709-723.

1140 Heikal, Y., Woll, M.P., Fox, T., Seaton, K., Levenson, R., and Kester, M. (2011). Neurotensin
 1141 receptor-1 inducible palmitoylation is required for efficient receptor-mediated
 1142 mitogenic-signaling within structured membrane microdomains. *Cancer biology &*
 1143 *therapy* 12, 427-435.

1144 Hemler, M.E. (2014). Tetraspanin proteins promote multiple cancer stages. *Nature reviews*
 1145 *Cancer* 14, 49-60.

1146 Hemsley, P.A. (2009). Protein S-acylation in plants (Review). *Molecular membrane biology*
 1147 26, 114-125.

1148 Hemsley, P.A. (2015). The importance of lipid modified proteins in plants. *New Phytologist*
 1149 205, 476-489.

1150 Hemsley, P.A., and Grierson, C.S. (2011). The ankyrin repeats and DHHC S-acyl transferase
 1151 domain of AKR1 act independently to regulate switching from vegetative to mating
 1152 states in yeast. *PloS one* 6, e28799.

1153 Hemsley, P.A., Kemp, A.C., and Grierson, C.S. (2005). The TIP GROWTH DEFECTIVE1 S-
 1154 acyl transferase regulates plant cell growth in Arabidopsis. *Plant Cell* 17, 2554-2563.

1155 Hemsley, P.A., Taylor, L., and Grierson, C.S. (2008). Assaying protein palmitoylation in plants.
 1156 *Plant Methods* 4, 1.

1157 Hemsley, P.A., Weimar, T., Lilley, K.S., Dupree, P., and Grierson, C.S. (2013). A proteomic
 1158 approach identifies many novel palmitoylated proteins in Arabidopsis. *New Phytologist*
 1159 197, 805-814.

1160 Hirschman, J.E., and Jenness, D.D. (1999). Dual lipid modification of the yeast Gγ subunit
 1161 Ste18p determines membrane localization of Gβγ. *Molecular and cellular biology* 19,
 1162 7705-7711.

- Hodson, N., Invergo, B., Rayner, J.C., and Choudhary, J.S. (2015). Palmitoylation and palmitoyl-transferases in Plasmodium parasites. *Biochemical Society Transactions* 43, 240-245.
- Hornemann, T. (2015). Palmitoylation and depalmitoylation defects. *Journal of inherited metabolic disease* 38, 179-186.
- Hou, H., Subramanian, K., Lagrassa, T.J., Markgraf, D., Dietrich, L.E., Urban, J., Decker, N., and Ungermann, C. (2005). The DHHC protein Pfa3 affects vacuole-associated palmitoylation of the fusion factor Vac8. *Proceedings of the National Academy of Sciences of the United States of America* 102, 17366-17371.
- Hu, L.L., Wan, S.B., Niu, S., Shi, X.H., Li, H.P., Cai, Y.D., and Chou, K.C. (2011). Prediction and analysis of protein palmitoylation sites. *Biochimie* 93, 489-496.
- Huang, K., Kang, M.H., Askew, C., Kang, R., Sanders, S.S., Wan, J., Davis, N.G., and Hayden, M.R. (2010). Palmitoylation and function of glial glutamate transporter-1 is reduced in the YAC128 mouse model of Huntington disease. *Neurobiology of disease* 40, 207-215.
- Huang, K., Sanders, S., Singaraja, R., Orban, P., Cijssouw, T., Arstikaitis, P., Yanai, A., Hayden, M.R., and El-Husseini, A. (2009). Neuronal palmitoyl acyl transferases exhibit distinct substrate specificity. *The FASEB Journal* 23, 2605-2615.
- Huang, K., Yanai, A., Kang, R., Arstikaitis, P., Singaraja, R.R., Metzler, M., Mullard, A., Haigh, B., Gauthier-Campbell, C., and Gutekunst, C.A. (2004). Huntingtin-interacting protein HIP14 is a palmitoyl transferase involved in palmitoylation and trafficking of multiple neuronal proteins. *Neuron* 44, 977-986.
- Hundt, M., Harada, Y., De Giorgio, L., Tanimura, N., Zhang, W., and Altman, A. (2009). Palmitoylation-dependent plasma membrane transport but lipid raft-independent signaling by linker for activation of T cells. *The Journal of Immunology* 183, 1685-1694.
- Hurst, C.H., and Hemsley, P.A. (2015). Current perspective on protein S-acylation in plants: more than just a fatty anchor? *Journal of experimental botany*, erv053.
- Ivaldi, C., Martin, B.R., Kieffer-Jaquinod, S., Chapel, A., Levade, T., Garin, J., and Journet, A. (2012). Proteomic analysis of S-acylated proteins in human B cells reveals palmitoylation of the immune regulators CD20 and CD23. *PLoS One* 7, e37187.
- Ivanov, S.S., and Roy, C. (2013). "Host lipidation: a mechanism for spatial regulation of Legionella effectors," in *Molecular Mechanisms in Legionella Pathogenesis*. Springer, 135-154.
- Jennings, B.C., Nadolski, M.J., Ling, Y., Baker, M.B., Harrison, M.L., Deschenes, R.J., and Linder, M.E. (2009). 2-Bromopalmitate and 2-(2-hydroxy-5-nitro-benzylidene)-benzo [b] thiophen-3-one inhibit DHHC-mediated palmitoylation in vitro. *Journal of lipid research* 50, 233-242.
- Ji, Y., Bachschmid, M.M., Costello, C.E., and Lin, C. (2016). S-to N-Palmitoyl Transfer During Proteomic Sample Preparation. *Journal of The American Society for Mass Spectrometry* 27, 677-685.
- Jones, T.L., Degtyarev, M.Y., and Backlund, P.S. (1997). The stoichiometry of Gαs palmitoylation in its basal and activated states. *Biochemistry* 36, 7185-7191.

- Kang, R., Wan, J., Arstikaitis, P., Takahashi, H., Huang, K., Bailey, A.O., Thompson, J.X., Roth, A.F., Drisdell, R.C., and Mastro, R. (2008). Neural palmitoyl-proteomics reveals dynamic synaptic palmitoylation. *Nature* 456, 904-909.
- Keith, D.J., Sanderson, J.L., Gibson, E.S., Woolfrey, K.M., Robertson, H.R., Olszewski, K., Kang, R., El-Husseini, A., and Dell'acqua, M.L. (2012). Palmitoylation of A-kinase anchoring protein 79/150 regulates dendritic endosomal targeting and synaptic plasticity mechanisms. *The Journal of Neuroscience* 32, 7119-7136.
- Keller, C.A., Yuan, X., Panzanelli, P., Martin, M.L., Alldred, M., Sassoè-Pognetto, M., and Lüscher, B. (2004). The $\gamma 2$ Subunit of GABA_A Receptors Is a Substrate for Palmitoylation by GODZ. *The Journal of Neuroscience* 24, 5881-5891.
- Kihara, A., Kurotsu, F., Sano, T., Iwaki, S., and Igarashi, Y. (2005). Long-chain base kinase Lcb4 is anchored to the membrane through its palmitoylation by Akr1. *Molecular and cellular biology* 25, 9189-9197.
- Kim, M.G., Da Cunha, L., Mcfall, A.J., Belkhadir, Y., Debroy, S., Dangl, J.L., and Mackey, D. (2005). Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in *Arabidopsis*. *Cell* 121, 749-759.
- Kleuss, C., and Krause, E. (2003). Gas is palmitoylated at the N-terminal glycine. *The EMBO journal* 22, 826-832.
- Konrad, S.S., Popp, C., Stratil, T.F., Jarsch, I.K., Thallmair, V., Folgmann, J., Marín, M., and Ott, T. (2014). S - acylation anchors remorin proteins to the plasma membrane but does not primarily determine their localization in membrane microdomains. *New Phytologist* 203, 758-769.
- Kostiuk, M.A., Corvi, M.M., Keller, B.O., Plummer, G., Prescher, J.A., Hangauer, M.J., Bertozzi, C.R., Rajaiah, G., Falck, J.R., and Berthiaume, L.G. (2008). Identification of palmitoylated mitochondrial proteins using a bio-orthogonal azido-palmitate analogue. *The FASEB Journal* 22, 721-732.
- Kumar, M., Wightman, R., Atanassov, I., Gupta, A., Hurst, C.H., Hemsley, P.A., and Turner, S. (2016). S-Acylation of the cellulose synthase complex is essential for its plasma membrane localization. *Science* 353, 166-169.
- Kumari, B., Kumar, R., and Kumar, M. (2014). PalmPred: an SVM based palmitoylation prediction method using sequence profile information. *PloS one* 9, e89246.
- Kümmel, D., Heinemann, U., and Veit, M. (2006). Unique self-palmitoylation activity of the transport protein particle component Bet3: a mechanism required for protein stability. *Proceedings of the National Academy of Sciences* 103, 12701-12706.
- Kurayoshi, M., Yamamoto, H., Izumi, S., and Kikuchi, A. (2007). Post-translational palmitoylation and glycosylation of Wnt-5a are necessary for its signalling. *Biochemical Journal* 402, 515-523.
- Lakkaraju, A.K., Abrami, L., Lemmin, T., Blaskovic, S., Kunz, B., Kihara, A., Dal Peraro, M., and Van Der Goot, F.G. (2012). Palmitoylated calnexin is a key component of the ribosome-translocon complex. *The EMBO journal* 31, 1823-1835.
- Lam, K.K., Davey, M., Sun, B., Roth, A.F., Davis, N.G., and Conibear, E. (2006). Palmitoylation by the DHHC protein Pfa4 regulates the ER exit of Chs3. *The Journal of cell biology* 174, 19-25.

1249 Lane, S.R., and Liu, Y. (1997). Characterization of the palmitoylation domain of SNAP-25.
1250 *Journal of neurochemistry* 69, 1864-1869.

1251 Lavy, M., Bracha-Drori, K., Sternberg, H., and Yalovsky, S. (2002). A cell-specific,
1252 prenylation-independent mechanism regulates targeting of type II RACs. *The Plant*
1253 *Cell* 14, 2431-2450.

1254 Lawrence, D.S., Zilfou, J.T., and Smith, C.D. (1999). Structure-activity studies of cerulenin
1255 analogues as protein palmitoylation inhibitors. *Journal of medicinal chemistry* 42,
1256 4932-4941.

1257 Leclercq, J., Ranty, B., Sanchez-Ballesta, M.-T., Li, Z., Jones, B., Jauneau, A., Pech, J.-C.,
1258 Latché, A., Ranjeva, R., and Bouzayen, M. (2005). Molecular and biochemical
1259 characterization of LeCRK1, a ripening-associated tomato CDPK-related kinase.
1260 *Journal of experimental botany* 56, 25-35.

1261 Lemonidis, K., Sanchez-Perez, M.C., and Chamberlain, L.H. (2015). Identification of a Novel
1262 Sequence Motif Recognized by the Ankyrin Repeat Domain of zDHHC17/13 S-
1263 Acyltransferases. *Journal of Biological Chemistry* 290, 21939-21950.

1264 Levental, I., Lingwood, D., Grzybek, M., Coskun, Ü., and Simons, K. (2010). Palmitoylation
1265 regulates raft affinity for the majority of integral raft proteins. *Proceedings of the*
1266 *National Academy of Sciences* 107, 22050-22054.

1267 Li, Y., Martin, B.R., Cravatt, B.F., and Hofmann, S.L. (2012). DHHC5 protein palmitoylates
1268 flotillin-2 and is rapidly degraded on induction of neuronal differentiation in cultured
1269 cells. *Journal of Biological Chemistry* 287, 523-530.

1270 Li, Y., Scott, R.J., Doughty, J., Grant, M., and Qi, B. (2015). Protein S-Acyltransferase 14: a
1271 specific role for palmitoylation in leaf senescence in Arabidopsis. *Plant physiology*, pp.
1272 00448.02015.

1273 Linder, M.E., and Deschenes, R.J. (2007). Palmitoylation: policing protein stability and traffic.
1274 *Nature reviews Molecular cell biology* 8, 74-84.

1275 Liu, J., Zhong, S., Guo, X., Hao, L., Wei, X., Huang, Q., Hou, Y., Shi, J., Wang, C., Gu, H.,
1276 and Qu, L.-J. (2013). Membrane-Bound RLCKs LIP1 and LIP2 Are Essential Male
1277 Factors Controlling Male-Female Attraction in Arabidopsis. *Current Biology* 23, 993-
1278 998.

1279 Liu, Y., Fisher, D.A., and Storm, D.R. (1993). Analysis of the palmitoylation and membrane
1280 targeting domain of neuromodulin (GAP-43) by site-specific mutagenesis.
1281 *Biochemistry* 32, 10714-10719.

1282 Lobo, S., Greentree, W.K., Linder, M.E., and Deschenes, R.J. (2002). Identification of a Ras
1283 palmitoyltransferase in *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*
1284 277, 41268-41273.

1285 Loisel, T.P., Adam, L., Hebert, T.E., and Bouvier, M. (1996). Agonist stimulation increases the
1286 turnover rate of β 2AR-bound palmitate and promotes receptor depalmitoylation.
1287 *Biochemistry* 35, 15923-15932.

1288 López-Barragán, M.J., Lemieux, J., Quiñones, M., Williamson, K.C., Molina-Cruz, A., Cui, K.,
1289 Barillas-Mury, C., Zhao, K., and Su, X.-Z. (2011). Directional gene expression and
1290 antisense transcripts in sexual and asexual stages of *Plasmodium falciparum*. *BMC*
1291 *genomics* 12, 1.

- Lu, D., Sun, H.-Q., Wang, H., Barylko, B., Fukata, Y., Fukata, M., Albanesi, J.P., and Yin, H.L. (2012). Phosphatidylinositol 4-kinase II α is palmitoylated by Golgi-localized palmitoyltransferases in cholesterol-dependent manner. *Journal of Biological Chemistry* 287, 21856-21865.
- Maciaszczyk-Dziubinska, E., Migocka, M., Wawrzycka, D., Markowska, K., and Wysocki, R. (2014). Multiple cysteine residues are necessary for sorting and transport activity of the arsenite permease Acr3p from *Saccharomyces cerevisiae*. *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1838, 747-755.
- Maeda, A., Okano, K., Park, P.S.-H., Lem, J., Crouch, R.K., Maeda, T., and Palczewski, K. (2010). Palmitoylation stabilizes unliganded rod opsin. *Proceedings of the National Academy of Sciences* 107, 8428-8433.
- Maisse, C., Rossin, A., Cahuzac, N., Paradisi, A., Klein, C., Haillot, M.-L., Hérincs, Z., Mehlen, P., and Hueber, A.-O. (2008). Lipid raft localization and palmitoylation: identification of two requirements for cell death induction by the tumor suppressors UNC5H. *Experimental cell research* 314, 2544-2552.
- Manandhar, S.P., Calle, E.N., and Gharakhanian, E. (2014). Distinct Palmitoylation Events at the Amino-terminal Conserved Cysteines of Env7 Direct Its Stability, Localization, and Vacuolar Fusion Regulation in *S. cerevisiae*. *Journal of Biological Chemistry* 289, 11431-11442.
- Manandhar, S.P., Ricarte, F., Cocca, S.M., and Gharakhanian, E. (2013). *Saccharomyces cerevisiae* Env7 is a novel serine/threonine kinase 16-related protein kinase and negatively regulates organelle fusion at the lysosomal vacuole. *Molecular and cellular biology* 33, 526-542.
- Marin, E.P., Derakhshan, B., Lam, T.T., Davalos, A., and Sessa, W.C. (2012). Endothelial cell palmitoylproteomic identifies novel lipid-modified targets and potential substrates for protein acyl transferases. *Circulation research* 110, 1336-1344.
- Martin, B.R., and Cravatt, B.F. (2009). Large-scale profiling of protein palmitoylation in mammalian cells. *Nature methods* 6, 135-138.
- Martin, B.R., Wang, C., Adibekian, A., Tully, S.E., and Cravatt, B.F. (2012). Global profiling of dynamic protein palmitoylation. *Nature methods* 9, 84-89.
- Mattoo, A.K., and Edelman, M. (1987). Intramembrane translocation and posttranslational palmitoylation of the chloroplast 32-kDa herbicide-binding protein. *Proceedings of the National Academy of Sciences* 84, 1497-1501.
- Mccormick, P.J., Dumaresq-Doiron, K., Pluviose, A.S., Pichette, V., Tosato, G., and Lefrancois, S. (2008). Palmitoylation controls recycling in lysosomal sorting and trafficking. *Traffic* 9, 1984-1997.
- Meiringer, C.T., and Ungermann, C. (2006). Probing protein palmitoylation at the yeast vacuole. *Methods* 40, 171-176.
- Merino, M.C., Zamponi, N., Vranich, C.V., Touz, M.C., and Ropolo, A.S. (2014). Identification of *Giardia lamblia* DHHC Proteins and the Role of Protein S-palmitoylation in the Encystation Process. *PLoS Negl Trop Dis* 8, e2997.
- Merrick, B.A., Dhungana, S., Williams, J.G., Aloor, J.J., Peddada, S., Tomer, K.B., and Fessler, M.B. (2011). Proteomic profiling of S-acylated macrophage proteins identifies a role

for palmitoylation in mitochondrial targeting of phospholipid scramblase 3. *Molecular & Cellular Proteomics* 10, M110. 006007.

Mill, P., Lee, A.W., Fukata, Y., Tsutsumi, R., Fukata, M., Keighren, M., Porter, R.M., Mckie, L., Smyth, I., and Jackson, I.J. (2009). Palmitoylation regulates epidermal homeostasis and hair follicle differentiation. *PLoS Genet* 5, e1000748.

Mitchell, D.A., Farh, L., Marshall, T.K., and Deschenes, R.J. (1994). A polybasic domain allows nonprenylated Ras proteins to function in *Saccharomyces cerevisiae*. *Journal of Biological Chemistry* 269, 21540-21546.

Mitchell, D.A., Vasudevan, A., Linder, M.E., and Deschenes, R.J. (2006). Thematic review series: lipid posttranslational modifications. Protein palmitoylation by a family of DHHC protein S-acyltransferases. *Journal of lipid research* 47, 1118-1127.

Miura, G.I., Buglino, J., Alvarado, D., Lemmon, M.A., Resh, M.D., and Treisman, J.E. (2006). Palmitoylation of the EGFR ligand Spitz by Rasp increases Spitz activity by restricting its diffusion. *Developmental cell* 10, 167-176.

Montoro, A.G., Ramirez, S.C., Quiroga, R., and Taubas, J.V. (2011). Specificity of transmembrane protein palmitoylation in yeast. *PLoS One* 6, e16969.

Motoki, K., Kume, H., Oda, A., Tamaoka, A., Hosaka, A., Kametani, F., and Araki, W. (2012). Neuronal β -amyloid generation is independent of lipid raft association of β -secretase BACE1: analysis with a palmitoylation-deficient mutant. *Brain and behavior* 2, 270-282.

Mukai, J., Dhilla, A., Drew, L.J., Stark, K.L., Cao, L., Macdermott, A.B., Karayiorgou, M., and Gogos, J.A. (2008). Palmitoylation-dependent neurodevelopmental deficits in a mouse model of 22q11 microdeletion. *Nature neuroscience* 11, 1302-1310.

Mukai, J., Liu, H., Burt, R.A., Swor, D.E., Lai, W.-S., Karayiorgou, M., and Gogos, J.A. (2004). Evidence that the gene encoding ZDHHC8 contributes to the risk of schizophrenia. *Nature genetics* 36, 725-731.

Mumby, S.M., Kleuss, C., and Gilman, A.G. (1994). Receptor regulation of G-protein palmitoylation. *Proceedings of the National Academy of Sciences* 91, 2800-2804.

Nagano, N., Ota, M., and Nishikawa, K. (1999). Strong hydrophobic nature of cysteine residues in proteins. *FEBS letters* 458, 69-71.

Nichols, C.B., Ost, K.S., Grogan, D.P., Pianalto, K., Hasan, S., and Alspaugh, J.A. (2015). Impact of protein palmitoylation on the virulence potential of *Cryptococcus neoformans*. *Eukaryotic cell* 14, 626-635.

Oddi, S., Dainese, E., Sandiford, S., Fezza, F., Lanuti, M., Chiurchiù, V., Totaro, A., Catanzaro, G., Barcaroli, D., and De Laurenzi, V. (2012). Effects of palmitoylation of Cys415 in helix 8 of the CB1 cannabinoid receptor on membrane localization and signalling. *British journal of pharmacology* 165, 2635-2651.

Oh, Y., Jeon, Y., Hong, G., Kim, I., Woo, H., and Jung, Y. (2012). Regulation in the targeting of TRAIL receptor 1 to cell surface via GODZ for TRAIL sensitivity in tumor cells. *Cell Death & Differentiation* 19, 1196-1207.

Ohno, Y., Kashio, A., Ogata, R., Ishitomi, A., Yamazaki, Y., and Kihara, A. (2012). Analysis of substrate specificity of human DHHC protein acyltransferases using a yeast expression system. *Molecular biology of the cell* 23, 4543-4551.

- Ohno, Y., Kihara, A., Sano, T., and Igarashi, Y. (2006). Intracellular localization and tissue-specific distribution of human and yeast DHHC cysteine-rich domain-containing proteins. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids* 1761, 474-483.
- Ohyama, T., Verstreken, P., Ly, C.V., Rosenmund, T., Rajan, A., Tien, A.-C., Haueter, C., Schulze, K.L., and Bellen, H.J. (2007). Huntingtin-interacting protein 14, a palmitoyl transferase required for exocytosis and targeting of CSP to synaptic vesicles. *The Journal of cell biology* 179, 1481-1496.
- Park, S., Patterson, E.E., Cobb, J., Audhya, A., Gartenberg, M.R., and Fox, C.A. (2011). Palmitoylation controls the dynamics of budding-yeast heterochromatin via the telomere-binding protein Rif1. *Proceedings of the National Academy of Sciences* 108, 14572-14577.
- Patterson, S.I., and Skene, J. (1995). Inhibition of dynamic protein palmitoylation in intact cells with tunicamycin. *Methods in enzymology* 250, 284.
- Pedro, M.P., Vilcaes, A.A., Tomatis, V.M., Oliveira, R.G., Gomez, G.A., and Daniotti, J.L. (2013). 2-Bromopalmitate reduces protein deacylation by inhibition of acyl-protein thioesterase enzymatic activities. *PloS one* 8, e75232.
- Pepinsky, R.B., Zeng, C., Wen, D., Rayhorn, P., Baker, D.P., Williams, K.P., Bixler, S.A., Ambrose, C.M., Garber, E.A., and Miatkowski, K. (1998). Identification of a palmitic acid-modified form of human Sonic hedgehog. *Journal of Biological Chemistry* 273, 14037-14045.
- Percherancier, Y., Planchenault, T., Valenzuela-Fernandez, A., Virelizier, J.-L., Arenzana-Seisdedos, F., and Bachelier, F. (2001). Palmitoylation-dependent control of degradation, life span, and membrane expression of the CCR5 receptor. *Journal of Biological Chemistry* 276, 31936-31944.
- Ponimaskin, E., Behn, H., Adarichev, V., Voyno-Yasenetskaya, T.A., Offermanns, S., and Schmidt, M.F. (2000). Acylation of Gα13 is important for its interaction with thrombin receptor, transforming activity and actin stress fiber formation. *FEBS letters* 478, 173-177.
- Ponimaskin, E., Dityateva, G., Ruonala, M.O., Fukata, M., Fukata, Y., Kobe, F., Wouters, F.S., Delling, M., Bredt, D.S., and Schachner, M. (2008). Fibroblast growth factor-regulated palmitoylation of the neural cell adhesion molecule determines neuronal morphogenesis. *The Journal of Neuroscience* 28, 8897-8907.
- Ponimaskin, E., Harteneck, C., Schultz, G., and Schmidt, M.F. (1998). A cysteine-11 to serine mutant of Gα12 impairs activation through the thrombin receptor. *FEBS letters* 429, 370-374.
- Qi, B., Doughty, J., and Hooley, R. (2013). A Golgi and tonoplast localized S-acyl transferase is involved in cell expansion, cell division, vascular patterning and fertility in Arabidopsis. *New Phytologist* 200, 444-456.
- Raíces, M., Ulloa, R.M., Macintosh, G.C., Crespi, M., and Téllez-Iñón, M.T. (2003). StCDPK1 is expressed in potato stolon tips and is induced by high sucrose concentration. *Journal of experimental botany* 54, 2589-2591.
- Raymond, F.L., Tarpey, P.S., Edkins, S., Tofts, C., O'meara, S., Teague, J., Butler, A., Stevens, C., Barthorpe, S., and Buck, G. (2007). Mutations in ZDHHC9, which encodes a

1422 palmitoyltransferase of NRAS and HRAS, cause X-linked mental retardation
 1423 associated with a Marfanoid habitus. *The American Journal of Human Genetics* 80,
 1424 982-987.

1425 Resh, M.D. (2006). Palmitoylation of ligands, receptors, and intracellular signaling molecules.
 1426 *Science Signaling* 2006, re14-re14.

1427 Robinson, L.J., Busconi, L., and Michel, T. (1995). Agonist-modulated palmitoylation of
 1428 endothelial nitric oxide synthase. *Journal of Biological Chemistry* 270, 995-998.

1429 Rocks, O., Gerauer, M., Vartak, N., Koch, S., Huang, Z.-P., Pechlivanis, M., Kuhlmann, J.,
 1430 Brunsveld, L., Chandra, A., and Ellinger, B. (2010). The palmitoylation machinery is
 1431 a spatially organizing system for peripheral membrane proteins. *Cell* 141, 458-471.

1432 Rocks, O., Peyker, A., Kahms, M., Verveer, P.J., Koerner, C., Lumbierres, M., Kuhlmann, J.,
 1433 Waldmann, H., Wittinghofer, A., and Bastiaens, P.I. (2005). An acylation cycle
 1434 regulates localization and activity of palmitoylated Ras isoforms. *Science* 307, 1746-
 1435 1752.

1436 Rossin, A., Derouet, M., Abdel-Sater, F., and Hueber, A. (2009). Palmitoylation of the TRAIL
 1437 receptor DR4 confers an efficient TRAIL-induced cell death signalling. *Biochemical*
 1438 *Journal* 419, 185-192.

1439 Roth, A.F., Feng, Y., Chen, L., and Davis, N.G. (2002). The yeast DHHC cysteine-rich domain
 1440 protein Akr1p is a palmitoyl transferase. *The Journal of cell biology* 159, 23-28.

1441 Roth, A.F., Papanayotou, I., and Davis, N.G. (2011). The yeast kinase Yck2 has a tripartite
 1442 palmitoylation signal. *Molecular biology of the cell* 22, 2702-2715.

1443 Roth, A.F., Wan, J., Bailey, A.O., Sun, B., Kuchar, J.A., Green, W.N., Phinney, B.S., Yates,
 1444 J.R., and Davis, N.G. (2006). Global analysis of protein palmitoylation in yeast. *Cell*
 1445 125, 1003-1013.

1446 Running, M.P. (2014). The role of lipid post-translational modification in plant developmental
 1447 processes. *Regulation of Cell Fate Determination in Plants*, 76.

1448 Saleem, A.N., Chen, Y.-H., Baek, H.J., Hsiao, Y.-W., Huang, H.-W., Kao, H.-J., Liu, K.-M.,
 1449 Shen, L.-F., Song, I.-W., and Tu, C.-P.D. (2010). Mice with alopecia, osteoporosis, and
 1450 systemic amyloidosis due to mutation in Zdhhc13, a gene coding for palmitoyl
 1451 acyltransferase. *PLoS Genet* 6, e1000985.

1452 Schmick, M., Kraemer, A., and Bastiaens, P.I. (2015). Ras moves to stay in place. *Trends in*
 1453 *cell biology* 25, 190-197.

1454 Schmidt, M.F., and Schlesinger, M.J. (1979). Fatty acid binding to vesicular stomatitis virus
 1455 glycoprotein: a new type of post-translational modification of the viral glycoprotein.
 1456 *Cell* 17, 813-819.

1457 Sharma, C., Rabinovitz, I., and Hemler, M.E. (2012). Palmitoylation by DHHC3 is critical for
 1458 the function, expression, and stability of integrin $\alpha 6 \beta 4$. *Cellular and Molecular Life*
 1459 *Sciences* 69, 2233-2244.

1460 Shchedrina, V.A., Everley, R.A., Zhang, Y., Gygi, S.P., Hatfield, D.L., and Gladyshev, V.N.
 1461 (2011). Selenoprotein K binds multiprotein complexes and is involved in the regulation
 1462 of endoplasmic reticulum homeostasis. *Journal of Biological Chemistry* 286, 42937-
 1463 42948.

- Shi, S.-P., Sun, X.-Y., Qiu, J.-D., Suo, S.-B., Chen, X., Huang, S.-Y., and Liang, R.-P. (2013). The prediction of palmitoylation site locations using a multiple feature extraction method. *Journal of Molecular Graphics and Modelling* 40, 125-130.
- Shmueli, A., Segal, M., Sapir, T., Tsutsumi, R., Noritake, J., Bar, A., Sapoznik, S., Fukata, Y., Orr, I., and Fukata, M. (2010). Ndel1 palmitoylation: a new mean to regulate cytoplasmic dynein activity. *The EMBO journal* 29, 107-119.
- Silvius, J.R., and L'heureux, F. (1994). Fluorometric evaluation of the affinities of isoprenylated peptides for lipid bilayers. *Biochemistry* 33, 3014-3022.
- Singaraja, R.R., Kang, M.H., Vaid, K., Sanders, S.S., Vilas, G.L., Arstikaitis, P., Coutinho, J., Drisdell, R.C., El-Husseini, A.E.D., and Green, W.N. (2009). Palmitoylation of ATP-binding cassette transporter A1 is essential for its trafficking and function. *Circulation research* 105, 138-147.
- Smotrys, J.E., and Linder, M.E. (2004). Palmitoylation of intracellular signaling proteins: regulation and function. *Annual review of biochemistry* 73, 559-587.
- Smotrys, J.E., Schoenfish, M.J., Stutz, M.A., and Linder, M.E. (2005). The vacuolar DHHC-CRD protein Pfa3p is a protein acyltransferase for Vac8p. *The Journal of cell biology* 170, 1091-1099.
- Song, I.-W., Li, W.-R., Chen, L.-Y., Shen, L.-F., Liu, K.-M., Yen, J.J., Chen, Y.-J., Chen, Y.-J., Kraus, V.B., and Wu, J.-Y. (2014). Palmitoyl acyltransferase, Zdhc13, facilitates bone mass acquisition by regulating postnatal epiphyseal development and endochondral ossification: a mouse model. *PloS one* 9, e92194.
- Song, J., and Dohlman, H.G. (1996). Partial constitutive activation of pheromone responses by a palmitoylation-site mutant of a G protein α subunit in yeast. *Biochemistry* 35, 14806-14817.
- Song, J., Hirschman, J., Gunn, K., and Dohlman, H.G. (1996). Regulation of membrane and subunit interactions by N-myristoylation of a G protein α subunit in yeast. *Journal of Biological Chemistry* 271, 20273-20283.
- Sorek, N., Poraty, L., Sternberg, H., Bar, E., Lewinsohn, E., and Yalovsky, S. (2007). Activation status-coupled transient S acylation determines membrane partitioning of a plant Rho-related GTPase. *Molecular and cellular biology* 27, 2144-2154.
- Soung, Y.H., and Chung, J. (2011). Curcumin inhibition of the functional interaction between integrin $\alpha 6 \beta 4$ and the epidermal growth factor receptor. *Molecular cancer therapeutics* 10, 883-891.
- Srivastava, V., Weber, J.R., Malm, E., Fouke, B.W., and Bulone, V. (2016). Proteomic analysis of a poplar cell suspension culture suggests a major role of protein S-acylation in diverse cellular processes. *Frontiers in plant science* 7.
- Swarthout, J.T., Lobo, S., Farh, L., Croke, M.R., Greentree, W.K., Deschenes, R.J., and Linder, M.E. (2005). DHHC9 and GCP16 constitute a human protein fatty acyltransferase with specificity for H- and N-Ras. *Journal of Biological Chemistry* 280, 31141-31148.
- Takada, R., Satomi, Y., Kurata, T., Ueno, N., Norioka, S., Kondoh, H., Takao, T., and Takada, S. (2006). Monounsaturated fatty acid modification of Wnt protein: its role in Wnt secretion. *Developmental cell* 11, 791-801.
- Takemoto-Kimura, S., Ageta-Ishihara, N., Nonaka, M., Adachi-Morishima, A., Mano, T., Okamura, M., Fujii, H., Fuse, T., Hoshino, M., and Suzuki, S. (2007). Regulation of

- dendritogenesis via a lipid-raft-associated Ca²⁺/calmodulin-dependent protein kinase
 CLICK-III/CaMKI γ . *Neuron* 54, 755-770.
- Termini, C.M., Cotter, M.L., Marjon, K.D., Buranda, T., Lidke, K.A., and Gillette, J.M. (2014).
 The membrane scaffold CD82 regulates cell adhesion by altering α 4 integrin stability
 and molecular density. *Molecular biology of the cell* 25, 1560-1573.
- Thaa, B., Levental, I., Herrmann, A., and Veit, M. (2011). Intrinsic membrane association of
 the cytoplasmic tail of influenza virus M2 protein and lateral membrane sorting
 regulated by cholesterol binding and palmitoylation. *Biochemical Journal* 437, 389-
 397.
- Tian, L., Mcclafferty, H., Jeffries, O., and Shipston, M.J. (2010). Multiple palmitoyltransferases
 are required for palmitoylation-dependent regulation of large conductance calcium-and
 voltage-activated potassium channels. *Journal of Biological Chemistry* 285, 23954-
 23962.
- Tomatis, V.M., Trenchi, A., Gomez, G.A., and Daniotti, J.L. (2010). Acyl-protein thioesterase
 2 catalyzes the deacylation of peripheral membrane-associated GAP-43. *PloS one* 5,
 e15045.
- Topinka, J.R., and Brecht, D.S. (1998). N-terminal palmitoylation of PSD-95 regulates
 association with cell membranes and interaction with K⁺ channel K v 1.4. *Neuron* 20,
 125-134.
- Toyoda, T., Sugimoto, H., and Yamashita, S. (1999). Sequence, expression in Escherichia coli,
 and characterization of lysophospholipase II. *Biochimica et Biophysica Acta (BBA)-
 Molecular and Cell Biology of Lipids* 1437, 182-193.
- Tsutsumi, R., Fukata, Y., Noritake, J., Iwanaga, T., Perez, F., and Fukata, M. (2009).
 Identification of G protein α subunit-palmitoylating enzyme. *Molecular and cellular
 biology* 29, 435-447.
- Ueda, T., Yamaguchi, M., Uchimiya, H., and Nakano, A. (2001). Ara6, a plant-unique novel
 type Rab GTPase, functions in the endocytic pathway of Arabidopsis thaliana. *The
 EMBO Journal* 20, 4730-4741.
- Valdez-Taubas, J., and Pelham, H. (2005). Swf1-dependent palmitoylation of the SNARE Tlg1
 prevents its ubiquitination and degradation. *The EMBO journal* 24, 2524-2532.
- Vesa, J., Hellsten, E., Verkruyse, L.A., Camp, L.A., Rapola, J., Santavuori, P., Hofmann, S.L.,
 and Peltonen, L. (1995). Mutations in the palmitoyl protein thioesterase gene causing
 infantile neuronal ceroid lipofuscinosis. *Nature*.
- Vetrivel, K.S., Meckler, X., Chen, Y., Nguyen, P.D., Seidah, N.G., Vassar, R., Wong, P.C.,
 Fukata, M., Kounnas, M.Z., and Thinakaran, G. (2009). Alzheimer disease A β
 production in the absence of S-palmitoylation-dependent targeting of BACE1 to lipid
 rafts. *Journal of Biological Chemistry* 284, 3793-3803.
- Wagner, R., Herwig, A., Azzouz, N., and Klenk, H.D. (2005). Acylation-mediated membrane
 anchoring of avian influenza virus hemagglutinin is essential for fusion pore formation
 and virus infectivity. *Journal of virology* 79, 6449-6458.
- Wang, X., Sutton, V.R., Peraza-Llanes, J.O., Yu, Z., Rosetta, R., Kou, Y.-C., Eble, T.N., Patel,
 A., Thaller, C., and Fang, P. (2007). Mutations in X-linked PORCN, a putative
 regulator of Wnt signaling, cause focal dermal hypoplasia. *Nature genetics* 39, 836-
 838.

- Webb, Y., Hermida-Matsumoto, L., and Resh, M.D. (2000). Inhibition of protein palmitoylation, raft localization, and T cell signaling by 2-bromopalmitate and polyunsaturated fatty acids. *Journal of Biological Chemistry* 275, 261-270.
- Wedegaertner, P.B., Chu, D., Wilson, P., Levis, M.J., and Bourne, H. (1993). Palmitoylation is required for signaling functions and membrane attachment of Gq alpha and Gs alpha. *Journal of Biological Chemistry* 268, 25001-25008.
- Wetzel, J., Herrmann, S., Swapna, L.S., Prusty, D., John Peter, A.T., Kono, M., Saini, S., Nellimarla, S., Wong, T.W., Wilcke, L., Ramsay, O., Cabrera, A., Biller, L., Heincke, D., Mossman, K., Spielmann, T., Ungermann, C., Parkinson, J., and Gilberger, T.W. (2015). The role of palmitoylation for protein recruitment to the inner membrane complex of the malaria parasite. *Journal of Biological Chemistry* 290, 1712-1728.
- Wolven, A., Okamura, H., Rosenblatt, Y., and Resh, M. (1997). Palmitoylation of p59fyn is reversible and sufficient for plasma membrane association. *Molecular Biology of the Cell* 8, 1159-1173.
- Xie, Y., Wolff, D.W., Wei, T., Wang, B., Deng, C., Kirui, J.K., Jiang, H., Qin, J., Abel, P.W., and Tu, Y. (2009). Breast cancer migration and invasion depend on proteasome degradation of regulator of G-protein signaling 4. *Cancer research* 69, 5743-5751.
- Yang, G., and Cynader, M.S. (2011). Palmitoyl acyltransferase zD17 mediates neuronal responses in acute ischemic brain injury by regulating JNK activation in a signaling module. *The Journal of Neuroscience* 31, 11980-11991.
- Yang, W., Di Vizio, D., Kirchner, M., Steen, H., and Freeman, M.R. (2010). Proteome scale characterization of human S-acylated proteins in lipid raft-enriched and non-raft membranes. *Molecular & Cellular Proteomics* 9, 54-70.
- Yeste-Velasco, M., Linder, M.E., and Lu, Y.-J. (2015). Protein S-palmitoylation and cancer. *Biochimica et Biophysica Acta (BBA)-Reviews on Cancer* 1856, 107-120.
- Yuan, X., Zhang, S., Sun, M., Liu, S., Qi, B., and Li, X. (2013). Putative DHHC-Cysteine-Rich Domain S-Acyltransferase in Plants. *PloS one* 8.
- Zeidman, R., Jackson, C.S., and Magee, A.I. (2009). Protein acyl thioesterases (Review). *Molecular membrane biology* 26, 32-41.
- Zeng, Q., Wang, X., and Running, M.P. (2007). Dual lipid modification of Arabidopsis Gγ-subunits is required for efficient plasma membrane targeting. *Plant physiology* 143, 1119-1131.
- Zhang, J., Planey, S.L., Ceballos, C., Stevens, S.M., Keay, S.K., and Zacharias, D.A. (2008). Identification of CKAP4/p63 as a major substrate of the palmitoyl acyltransferase DHHC2, a putative tumor suppressor, using a novel proteomics method. *Molecular & Cellular Proteomics* 7, 1378-1388.
- Zhang, M.M., Wu, P.-Y.J., Kelly, F.D., Nurse, P., and Hang, H.C. (2013). Quantitative control of protein S-palmitoylation regulates meiotic entry in fission yeast. *PLoS Biol* 11, e1001597.
- Zhao, L., Lobo, S., Dong, X., Ault, A.D., and Deschenes, R.J. (2002). Erf4p and Erf2p form an endoplasmic reticulum-associated complex involved in the plasma membrane localization of yeast Ras proteins. *Journal of Biological Chemistry* 277, 49352-49359.

- 1594 Zhou, B., Liu, L., Reddivari, M., and Zhang, X.A. (2004). The palmitoylation of metastasis
1595 suppressor KAI1/CD82 is important for its motility-and invasiveness-inhibitory
1596 activity. *Cancer research* 64, 7455-7463.
- 1597 Zhou, F., Xue, Y., Yao, X., and Xu, Y. (2006). CSS-Palm: palmitoylation site prediction with
1598 a clustering and scoring strategy (CSS). *Bioinformatics* 22, 894-896.
- 1599 Zhou, L.Z., Li, S., Feng, Q.-N., Zhang, Y.-L., Zhao, X., Zeng, Y.-L., Wang, H., Jiang, L., and
1600 Zhang, Y. (2013). PROTEIN S-ACYL TRANSFERASE10 Is Critical for
1601 Development and Salt Tolerance in Arabidopsis. *The Plant Cell* 25, 1093-1107
- 1602 Zurcher, T., Luo, G., and Palese, P. (1994). Mutations at palmitoylation sites of the influenza
1603 virus hemagglutinin affect virus formation. *Journal of virology* 68, 5748-5754.
- 1604

Provisional

FIGURE LEGENDS

FIGURE 1 | Formulae of N-myristoylation, S-acylation and prenylation. For N-myristoylation, a 14-carbon myristoyl group is covalently attached by an amide bond to the alpha-amino group of an N-terminal glycine (G, in red); S-acylation is the attachment of a 16-carbon palmitate to cysteine residue (C, in red) via thioester bond; and Prenylation makes a 15-carbon farnesyl link to the CaaX cysteine residue in C-termini.

FIGURE 2 | Topology structure and conserved domains of PATs. Most PATs have 4 transmembrane domains (TMDs, blue columns) and their N- and C-termini are in the cytoplasm. A highly conserved catalytic DHHC-CRD (aspartate-histidine-histidine-cysteine cysteine rich domain) resides between the 2nd- and 3rd-TMDs. The majority of PATs also have the DPG (aspartate-proline-glycine), TTxE (threonine-threonine-any amino acid-glutamic acid) and PaCCT (Palmitoyltransferase Conserved C-Terminus) domains, and all of them are cytosolic.

Provisional




Figure 02.JPEG

